



JAN 17, 2024

Fixed RNA - GEM Recovery - Sequencing

Forked from [Fixed RNA - FFPE Resection Tissue \(gentleMACS dissociation\)](#)

Ksenija
Sabic¹

¹Icahn School of Medicine at Mount Sinai



Ksenija Sabic

ABSTRACT

This protocol is a continuation of CG000527 (Step 3 and onward).

IMAGE ATTRIBUTION

10x Genomics

GUIDELINES

Please review and consult the full 10x Genomics protocols prior to starting and at any point during the procedure if needed.

OPEN ACCESS



DOI:

dx.doi.org/10.17504/protocols.io.3byl4q8ejvo5/v1

Protocol Citation: Ksenija Sabic 2024. Fixed RNA - GEM Recovery - Sequencing .

protocols.io

<https://dx.doi.org/10.17504/protocols.io.3byl4q8ejvo5/v1>

MANUSCRIPT CITATION:

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working


Created: Dec 26, 2023

Last Modified: Jan 17, 2024

PROTOCOL integer ID:
92741

MATERIALS


From 10x Genomics:

-  Chromium Fixed RNA Kit, Human Transcriptome, 4 rxns x 4 BC 10x
Genomics Catalog #1000475






OR

-  Chromium Fixed RNA Kit, Human Transcriptome, 4 rxns x 16 BC 10x
Genomics Catalog #1000476
-  Dual Index Kit TS Set A, 96 rxns 10x
Genomics Catalog #1000251

From Agilent:

-  Bioanalyzer High Sensitivity DNA Kit Agilent
Technologies Catalog #5067-4626

Miscellaneous:

-  SPRIselect Beckman Coulter Catalog #B23318
-  10% Tween 20 solution
Teknova Catalog #T0710 or similar
-  Ethyl alcohol, 200 proof, anhydrous, $\geq 99.5\%$ Merck MilliporeSigma (Sigma-Aldrich) Catalog #459836
- or similar
-  Nuclease-Free Water (not DEPC-Treated) Thermo Fisher
Scientific Catalog #AM9937
- or similar
-  Buffer
EB Qiagen Catalog #19086 or similar

BEFORE START INSTRUCTIONS

Prepare daily fresh 80% Ethanol for wash steps.

GEM Recovery and Pre-Amplification


3d 0h 9m

1 Reagent Preparation (~15 mins):

1.1 Thaw **Reducing Agent B** at  Room temperature, vortex, verify no precipitate, centrifuge briefly.

1.2 Thaw **Pre-Amp Primers B** at  Room temperature, vortex, centrifuge briefly.

1.3 Keep **Amp Mix**  On ice. Vortex and centrifuge briefly.

1.4 Prepare fresh 80% Ethanol ( 2.5 mL for 4 GEM reactions).

2 Add  125 μ L Recovery Agent to each sample at room temperature.



DO NOT pipette mix or vortex the biphasic mixture.


3 Firmly secure the cap on the tube strip, ensuring that no liquid is trapped between the cap and the tube rim. Mix by inverting the capped tube strip 5x.

DO NOT invert without firmly securing the caps.

4 Wait 2 min.

The resulting biphasic mixture contains Recovery Agent/Partitioning Oil (pink) and aqueous phase (translucent/opaque). A smaller aqueous phase volume indicates a clog during GEM generation.


5 Centrifuge briefly.

6 Slowly remove and discard  125 µL Recovery Agent/Partitioning Oil (pink) from the bottom of the tube. *DO NOT aspirate any aqueous sample.*

7 Proceed directly to Pre-Amplification PCR. No cleanup step is required.

8 Prepare Pre-Amplification Mix on ice. Vortex and centrifuge briefly.

A	B	C	D
Pre-Amp Mix (add reagents in order listed)	PN	1X (µl)	4X + 10% (µl)
Amp Mix	2000103	25.0	110.0
Pre-Amp Primers B	2000529	10.0	44.0
Total		35.0	154.00

9 Add  35 µL Pre-Amplification Mix to aqueous sample from step 6.

10 Cap firmly and invert 8x to mix. Centrifuge briefly.

11 Incubate in a thermal cycler with the following protocol:



A	B	C
Step	Temperature	Time (hh:mm:ss)
1	98C	00:03:00
2	98C	00:00:15
3	63C	00:00:20
4	72C	00:01:00
5	Go to Step 2, 7x (total 8 cycles)	

A	B	C
6	72C	00:01:00
7	4C	Hold

Saved as 'pre-amp' under Fixed RNA folder.

Lid Temperature: 105C

Reaction Volume: 100µl

Run Time: ~30-45 min

Store at 4°C for up to 72 h or -20°C for ≤1 week, or proceed to the next step.

12 Prepare Elution Solution. Vortex and centrifuge briefly.

A	B	C
Elution Solution (add reagents in order listed)	PN	1000µl
Buffer EB		980
10% Tween 20		10
Reducing Agent B	2000087	10
Total		1000







13 Centrifuge the sample (PCR product) for 00:00:30 sec in a microcentrifuge and transfer 70 30s of the upper layer to a new tube.


Presence of a cloudy precipitate at the interface between phases is normal. Avoid transferring the precipitate when transferring 70 µl at this step.

14 Vortex to resuspend the SPRIselect reagent. Add 126 µL **SPRIselect reagent** (1.8X) to each sample and pipette mix 15x (pipette set to 180 µl).

15 Incubate 00:05:00 min at Room temperature 5m.

16 Place on the magnet (high position) until the solution clears.

- 17 Remove the supernatant. ***DO NOT discard any beads.***
- 18 With the tube still in the magnet, add  200 µL **80% ethanol** to the pellet. Wait  00:00:30 sec. **30s**
- 18.1 Remove the ethanol.
- 18.2 Repeat steps 18 and 18.1 for a total of 2 washes.
- 19 Centrifuge briefly and place on the magnet (low position).
- 20 Remove any remaining ethanol. ***DO NOT let the sample dry to ensure maximum elution efficiency.***
- 21 Remove from the magnet. Add  101 µL **Elution Solution** (from step 12). Wait  00:01:00 min **1m**
before resuspending. Pipette mix 15x.
- 22 Incubate  00:02:00 min at  Room temperature **2m**
- 23 Place the tube strip on the magnet (high position) until the solution clears.

24 Transfer  100 µL sample to a new tube strip.

25 Store at  4 °C for ≤  72:00:00 hours or at  -20 °C for ≤4 weeks, or proceed to the next step, 






Library Construction

7m 30s

26 Reagent Preparation (~10 min):

26.1 Thaw **Dual Index Plate TS Set A** at  Room temperature .

26.2 Keep **Amp Mix**  On ice . Vortex and centrifuge briefly.

26.3  If planning to assess quality and concentration of library immediately after completion, thaw **Agilent Bioanalyzer High Sensitivity Kit** at  Room temperature , keeping prepared gel and gel-dye away from light sources. Consult the following protocol for details:

Protocol



NAME

Bioanalyzer High Sensitivity DNA Kit

CREATED BY


Ksenija Sabic

PREVIEW

27 Choose the appropriate sample index sets to ensure that no sample indices overlap in a multiplexed sequencing run. Record the 10x sample index name (PN-3000511 Dual Index Plate TS Set A well ID) used.

28 Prepare Sample Index PCR Mix on ice:

A	B	C	D	E
Sample Index PCR (add reagents in the order listed)	PN	1X (μl)	1X + 10% (μl)	4X + 10% (μl)
Amp Mix	2000103	50	55	220
Nuclease-free water		10	11	44
Total		60	66	264

29 Transfer **ONLY**  20 μL of sample from the step 24 to a new tube strip.

30 Add  60 μL **Sample Index PCR Mix** to 20 μl sample.

31 Add  20 μL of an individual **Dual Index TS Set A** to each sample. Pipette mix 5x (pipette set to 90 μl). Centrifuge briefly.

32 Incubate in a thermal cycler with the following protocol:

A	B	C
Step	Temperature	Time (hh:mm:ss)
1	98C	00:00:45
2	98C	00:00:20
3	54C	00:00:30
4	72C	00:00:20

A	B	C
5	Go to step 2, see table below for total # of cycles	
6	72C	00:01:00
7	4C	Hold

Saved as 'Sample Index PCR' under Fixed RNA folder.

Lid Temperature: 105C

Reaction Volume: 100µl

Run Time: ~25 - 40 min

32.1

A	B	C	D	E
Targeted Cell Recovery	Total Cycles*			
	for Cell Lines	for PBMCs & Nuclei	for Cells from Fixed & Dissociated Tissues**	for Cells from FFPE Tissue Sections
500 - 2,000	11	15	14 - 15	16
2,000 - 4,000	10	14	13 - 14	15
4,000 - 7,000	9	13	12 - 13	14
7,000 - 12,000	8	12	11 - 12	13
12,000 - 25,000	7	11	10 - 11	12
25,000 - 50,000	6	10	9 - 10	11
50,000 - 128,000	5	9	8 - 9	10

**Optimization of cycle number may be needed based on the total RNA content of the sample. The ideal target library concentration is 50 - 200 nM. However, if the concentration is between 10-50 nM or between 200-500 nM and if the libraries do not contain low or high molecular weight peaks, sequencing can still be performed. If optimization is needed, additional Amp Mix can be obtained using the Fixed RNA Feature Barcode Kit (PN-1000419). For dissociated tumor cells, cycle numbers for cell lines can be used as a starting point. For dissociated primary cells, cycle numbers for PBMCs can be used as a starting point.*

***For cells derived from the fixed and dissociated tissue samples, the cycle number will depend on the RNA expression level of the tissue and on overall quality of the tissue prior to fixation. Additional optimization may be required.*

33 Store at 4 °C for ≤72 h, or proceed to the next step.



34 Vortex to resuspend the SPRIselect reagent. Add 100 µL **SPRIselect Reagent** (1.0X) to each sample. Pipette mix 15x (pipette set to 180 µl).

35 Incubate 00:05:00 min at Room temperature .

5m

36 Place on the magnet (high position) until the solution clears.

37 Remove the supernatant. **DO NOT discard any beads.**

38 With the tube still in the magnet, add 200 µL **80% ethanol** to the pellet. Wait 00:00:30 sec. 30s

38.1 Remove the ethanol.

38.2 Repeat steps 38 and 38.1 for a total of 2 washes.


39 Centrifuge briefly and place on the magnet (low position).

40 Remove any remaining ethanol. ***DO NOT let the sample dry to ensure maximum elution efficiency.***



41 Remove from the magnet. Add  41 µL **Buffer EB**. Pipette mix 15x.

42 Incubate  00:02:00 min at  Room temperature . 2m


43 Place on the magnet (low position) until the solution clears.

44 Transfer  40 µL to a new labeled DNA LoBind tube.

44.1 Make a 1:80 dilution of the library for quantification.

45 Store at  4 °C for up to 72 h,  -20 °C or at  -80 °C for long-term storage.



46 Run  1 µL **sample at 1:80 dilution** on an **Agilent Bioanalyzer High Sensitivity chip**. Select the region between 150-300 bp to determine average size of the library. Consult the following protocol:

Protocol



NAME


Bioanalyzer High Sensitivity DNA Kit

CREATED BY

Ksenija Sabic

PREVIEW

46.1 See image below for representative trace:



Sequencing

47 Consult the following protocol to send the libraries for sequencing:

Protocol



NAME

Sequencing 10x Single Cell Libraries (UMGC Workflow)

CREATED BY

Ksenija Sabic

PREVIEW