



VERSION 1

MAY 24, 2023

OPEN  ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.e6nvwjk7dlmk/v1

Protocol Citation: Diana Rose E Ranoa 2023. SRB assay for measuring target cell killing. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.e6nvwjk7dlmk/v1> Version created by [Diana Rose E Ranoa](#)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited


Protocol status: Working
We use this protocol and it's working

Created: Oct 27, 2022**Last Modified:** May 24, 2023**PROTOCOL integer ID:**
71933

SRB assay for measuring target cell killing V.1

 In 2 collections

Diana Rose E
Ranoa¹

¹University of Illinois at Urbana-Champaign

Diana Rose E Ranoa
University of Illinois at Urbana-Champaign

DISCLAIMER

Protocol adapted from:
Vichai V, Kirtikara K. Sulforhodamine B colorimetric assay for cytotoxicity screening. *Nature Protocols* **2006**;1(3):1112-6 doi 10.1038/nprot.2006.179.

ABSTRACT

The Sulforhodamine B colorimetric assay was used to measure effector CAR T cell killing of their target tumor cells.

- 1 Perform co-culture experiment of activated CAR T cells and target tumor cells at various tumor:effector ratio in a 96-well plate for 24 hours at 37'C, 5% CO2.
- 2 Remove cell culture supernatant 24 hours after co-culture assay.
- 3 Using a multichannel pipet, fix adherent tumor cells on the 96-well plate with 100uL cold 10% (w/v) trichloroacetic acid (TCA) to each well, and incubate at 4'C for 1hr.
- 4 Wash four times by dipping entire plate in a basin with slow-running tap water. Tap plates on paper towels to remove excess water, and allow to dry at room temperature.
- 5 Add 100uL of 0.057% (w/v) Sulforhodamine B (SRB) solution to each well and incubate at room temperature for 30 minutes.
- 6 Immediately wash plates four times in a basin containing 1% (v/v) acetic acid to remove unbound dye.
- 7 After the plates have dried completely, add 200uL of 10mM Tris base solution (pH 10.5) to each well and incubate for 5 minutes with shaking.
- 8 Read plates in a microplate reader at A510nm.
- 9 Calculate the percentage of cell growth inhibition using the formula below:
$$\% \text{ cells killed} = 100 - [\text{mean OD}(\text{co-cultured sample}) / \text{mean OD}(\text{tumor cell only})] \times 100$$

