

Oct 24, 2020

pYCR cloning strategy

Andreea S¹¹University of Groningen

Other dx.doi.org/10.17504/protocols.io.bnvxme7n

iGEM Groningen 2020

a.stan.6

ABSTRACT

Introducing NLP14a in the genome of *B. mycoides*: In order to create plasmid pYCR-gamyNLP, the backbone of Pycr will be digested with the PCR product of gamy_Fw and gamy_Rv to produce the pYCR_gamy (CRISPR vector containing the gRNA). The NLP14a sequence will be ordered from Twist Biosciences And PCR-amplified with the primers NLP_Fw and NLP_Rv. The resulting PCR product, as well as the pYCR_gamy vector will be digested with SfiI. The digested mixtures will be ligated using T4 ligase.

Kill switch cloning: The suggested kill switch mechanism is based on Trp auxotrophic strains as well as a toxin-antitoxin mechanism. The toxin will be introduced in the genome using crispr in place of trpE. The antitoxin will be maintained as a cytoplasmic plasmid (pAD-YqcF) and expressed only in the presence of solanine. Unfortunately, to the best of our knowledge, no operator that binds solanine of *B. mycoides* has been described in literature. If it is proven that no such operator is present, we suggest on only keeping the tryptophan dependence strategy.

In order to create plasmid pYCR-gtrpEYqcG plasmid, the backbone of Pycr will be digested with BsaI and ligated with the annealing product of gtrpe_Fw and gtrpe_Rv to produce the pYCR_gtrp (CRISPR vector containing the gRNA). Synthetic dna for *yqcG* will be ordered from Twist Biosciences and PCR-amplified with the primers Yqcg_Fw and Yqcg_Rv. The resulting PCR product, as well as the pYCR_gtrpE vector will be digested with SfiI. The digested mixtures will be ligated using T4 ligase

Protocol was adapted after a paper of Yi et al., 2018

1. Yi, Y., Li, Z., Song, C. & Kuipers, O. P. Exploring plant-microbe interactions of the rhizobacteria *Bacillus subtilis* and *Bacillus mycoides* by use of the CRISPR-Cas9 system. *Environ. Microbiol.* 20, 4245–4260 (2018).

DOI

dx.doi.org/10.17504/protocols.io.bnvxme7n

PROTOCOL CITATION

Andreea S 2020. pYCR cloning strategy . protocols.io
<https://dx.doi.org/10.17504/protocols.io.bnvxme7n>

LICENSE

This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Oct 23, 2020

LAST MODIFIED

Oct 24, 2020

PROTOCOL INTEGER ID

ABSTRACT

Introducing NLP14a in the genome of *B. mycoides*: In order to create plasmid pYCR-gamyNLP, the backbone of Pycr will be digested with the PCR product of gamy_Fw and gamy_Rv to produce the pYCR_gamy (CRISPR vector containing the gRNA). The NLP14a sequence will be ordered from Twist Biosciences And PCR-amplified with the primers NLP_Fw and NLP_Rv. The resulting PCR product, as well as the pYCR_gamy vector will be digested with Sfil. The digested mixtures will be ligated using T4 ligase.







Kill switch cloning: The suggested kill switch mechanism is based on Trp auxotrophic strains as well as a toxin-antitoxin mechanism. The toxin will be introduced in the genome using crispr in place of trpE. The antitoxin will be maintained as a cytoplasmic plasmid (pAD-YqcF) and expressed only in the presence of solanine. Unfortunately, to the best of our knowledge, no operator that binds solanine of *B. mycoides* has been described in literature. If it is proven that no such operator is present, we suggest on only keeping the tryptophan dependence strategy.




In order to create plasmid pYCR-gtrpEYqcG plasmid, the backbone of Pycr will be digested with BsaI and ligated with the annealing product of gtrpe_Fw and gtrpe_Rv to produce the pYCR_gtrp (CRISPR vector containing the gRNA). Synthetic dna for *yqcG* will be ordered from Twist Biosciences and PCR-amplified with the primers Yqcg_Fw and Yqcg_Rv. The resulting PCR product, as well as the pYCR_gtrpE vector will be digested with Sfil. The digested mixtures will be ligated using T4 ligase

Protocol was adapted after a paper of Yi et al., 2018

1. Yi, Y., Li, Z., Song, C. & Kuipers, O. P. Exploring plant-microbe interactions of the rhizobacteria *Bacillus subtilis* and *Bacillus mycoides* by use of the CRISPR-Cas9 system. *Environ. Microbiol.* 20, 4245–4260 (2018).

Clone sgRNA sequence into pYCR

- 1 Design sgRNA spacer sequence (~20 nts) using "Benchling" and choose *B. mycoides* M2E_15 genome. Choose the gRNA with the highest on-target score and the lowest off-target score.
- 2 Order the 2 complementary oligos flanked by overhang containing BsaI restriction site
- 3 Anneal oligos by mixing the following components:
 -  **70 µl** Nuclease Free water
 -  **10 µl** Oligo DNA annealing buffer 10x (**100 Millimolar (mM)** Tris **pH8** , **500 Millimolar (mM)** NaCl, **10 Millimolar (mM)** EDTA)
 -  **10 µl** DNA oligos A and B (**10 Micromolar (µM)** each)
- 4 Mix well and incubate for  **00:05:00** at  **90 °C** 5m
- 5 Cool slowly to room temperature (aprox.  **01:00:00**) by removing the heat block from the apparatus 1h



- 6 Run agarose gel (1%) to check the annealing product
- 7 Digest pYCR with BsaI and inactivate the enzyme by PCR clean up
- 8 Ligate the annealing product with the precut pYCR:
Mlx:
 -  **10 µl** of annealing product
 -  **60 ng** of digested pYCR
 -  **1 µl** T4 ligase
- 9 Transform the ligation mixture into E. coli MC1061 using the heat-shock method.
Grow the cells on LB agar plates + Erythromycin (**0.1 mg/ml**)
- 10 To select the transformants perform a colony PCR with Fw primer binding upstream of PvuII and Rv primer downstream SfiI cloning sites.



Successful transformation will have a size of 427 pb

- 11 Miniprep and sequence the constructs

Clone flanking regions into pYCR containing sgRNA

- 12 Select the repair fragment of choice:
 - for knock-out: PCR amplify approx. 1 kbp upstream and downstream of the sgRNA targeted region
 - for knock-in: sequence of the desired insertion sequence flanked by SfiI restriction sites compatible with the backbone of pYCR
- 13 Digest the pYCR(+sgRNA) and the repair fragments with SfiI
Inactivate SfiI by PCR clean up
- 14 Ligate the components using T4 ligase 20m
Heat inactivate the ligase (incubate  **00:20:00**  **60 °C**)
- 15 Perform SmaI digestion to get rid of the original pYCR

- 16 Transform the ligation mixture into *E. coli* MC1061 using the heat-shock method. Grow the cells on LB agar plates + erythromycin
- 17 To select the transformants perform a colony PCR with Fw primer binding upstream of P_{vanP} and Rv primer downstream S_{fil} cloning sites.
- 18 Miniprep and sequence the constructs

Transform final construct in *B. mycoides* M2E_15




- 19 Transform using an electroporation protocol using  5 µl plasmid DNA

**Cloning of *Bacillus mycoides***
by a.stan.6

PREVIEW

RUN



- 19.1 Pick 1 colony of *B. mycoides* M2E_15 and inoculate it in BHIS
- 19.2 When OD_{600nm} reached 0.85 add 2% glycine and 2% threonine in order to weaken the cell wall.
- 19.3 Grow the cells overnight at  200 rpm, 30°C
- 19.4 Dilute the overnight culture 50 times in LBSP medium until the OD_{600nm} reaches 0.65
- 19.5 Collect the cells by  4000 x g, 4°C, 00:10:00 . Discard supernatant
- 19.6 Wash cell pellet with pre-chilled electroporation buffer (10% glycerol, 0.25 M sorbitol, 0.25M trehalose) x4
- 19.7 Suspended in 1 ml electroporation buffer (10% glycerol, 0.25 M sorbitol, 0.25M trehalose).
- 19.8 Snap-freeze the electrocompetent cells in liquid nitrogen and store at  -80 °C

- 19.9 For the PCR mix pipette:
- **5 µl** primers mixture Fw + Rv (5 uM) (see table "Primers")
 - **10 µl** HF buffer
 - **1 µl** dNTP (10mM)
 - DNA template (0.1 ng)
 - **0.25 µl** phusion polymerase
 - MiliQ to **50 µl**

19.10 PCR reaction is performed with the following protocol:

Temperature (*C)	Time (mm:ss)
98	05:00
98	10:00
Variable	00:30
72	00:30
repeat above steps	30x
72	10:00
11	Infinite

- 19.11 Load **5 µl** of sample plus **1 µl** of staining solution 6x. Check the length of the band (bp).
Purify sample : PCR cleanup/ gel extraction according to manufacturer.
- 19.12 Digest the polymerized insert and the desired vector with 2U of appropriate restriction enzyme. See table "Primers" in order to choose the enzyme.
- 19.13 Digestion mixture (**20 µl**):
- Sfil 2 U
 - Plasmid (**50 ng**)
 - Insert (**20 ng**)
 - **2 µl** CutSmart 10x (or appropriate buffer)
 - MiliQ to **20 µl**
- 19.14 Incubate **01:00:00** at the temperature recommended by the manufacturer of the restriction enzyme
- 19.15 Inactivate the restriction enzyme by incubating 20 min **65 °C**
**Sfil can't be heat inactivated so in order to inactivate it a PCR clean up is necessary

- 19.16 Ligate the digested insert and vector.
Ligation mixture :
-10 ul digestion product
-1 ul T4
-2 ul T4 ligase buffer
-7 ul MQ
- 19.17 Incubate 🕒 **02:00:00** 🌡 **Room temperature**
- 19.18 Thaw on ice 🧊 **100 µl** of electrocompetent cells
- 19.19 Add 🧊 **2 µg** of plasmid and the aliquot of electrocompetent cells to ice-cold electroporation cuvettes
- 19.20 Electroporate 25 uF, 10 kV/cm, 200 Ohms
- 19.21 Add 1 ml of BHIS and incubate 🌀 **100 rpm, 30°C, 05:00:00** for recovery
- 19.22 Plate on LB + Agar (1.5%) + 100 µg/ml spectinomycin.
For crispr cloning, to activate cas9 expression, add 0.2% mannose.
- 19.23 Incubate at 🌡 **30 °C** 🕒 **Overnight**
- 19.24 Randomly pick potential mutants and test by colony PCR (for primers use the ones that flank the region of insertion sites).
- 19.25 After selecting colonies that show the expected band size, purify the desired plasmid (miniprep) and transform into B. mycoides.
- 20 Plate on LB agar containing erythromycin
Grow at 🌡 **28 °C** overnight
- 21 Pick colonies and test them by colony PCR
If the expected band size is present, purify the plasmid and sequence

- 22 Grow the colonies with the expected sequence on liquid LB + erythromycin + manose
Incubate 🌀 **200 rpm, 28°C, 12:00:00**
- 23 Next day pick colonies in BHI medium (without antibiotic) and grow at 🌡 **37 °C** overnight
- 24 Dilute the liquid cultures and plate them on LN (no antibiotic)
grow overnight st 🌡 **37 °C**