



Aug 14, 2020

Universal Immunoblot analysis for investigating Protein-LAG (SpLAG)-binding to mammalian and avian immunoglobulins.

Angel A Justiz-Vaillant¹¹University of the West Indies St. Augustine**1** Works for me dx.doi.org/10.17504/protocols.io.bjsdkna6[University of the West Indies](#) angel.vaillant@sta.uwi.eduAngel Justiz-Vaillant
University of the West Indies St. Augustine

ABSTRACT

A protein that combines the binding capacity of SpA, SpG and SpL is not commercially available. It was created in my laboratory by combining these 3 immunoglobulin-binding proteins to horseradish peroxidase by the periodate method [1]. However, a mixture of SpA, SpG and SpL which are commercially available can have the same effect as universal reagent in immunodetection.

1. Vaillant AJ, McFarlane-Anderson N, Wisdom B, Mohammed W, Vuma S, et al. (2013) Immunoglobulin-binding Bacterial Proteins (IBP) Conjugates and their Reactivity with Immunoglobulin in Enzyme-Linked Immunosorbent Assays (ELISA). J Anal Bioanal Tech 4: 175. doi:10.4172/2155-9872.1000175

DOI

dx.doi.org/10.17504/protocols.io.bjsdkna6

PROTOCOL CITATION

Angel A Justiz-Vaillant 2020. Universal Immunoblot analysis for investigating Protein-LAG (SpLAG)-binding to mammalian and avian immunoglobulins.. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bjsdkna6>

LICENSE

This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Aug 14, 2020

LAST MODIFIED

Aug 14, 2020

PROTOCOL INTEGER ID

40485

- 1 Aliquots of egg yolks, sera or 2 µg/µl of purified immunoglobulins from birds, laboratory, wild, farm animals and pets are applied to the gels of SDS-PAGE as described elsewhere.
- 2 Gels are transferred to nitrocellulose membranes (Immobilon-Nc, pore size 0.45 µm, Sigma-Aldrich Co, St Louis, Missouri) during 71 minutes at 40 mAmps using a semi-dry electroblotter, HEP-1 Model, Owl Scientific Inc.

- 3 The running buffer contains 25 mM Tris, 192 mM glycine pH 8.3 and 20% methanol.
- 4 The nitrocellulose membranes are blocked overnight in 10% non-fat skim milk in PBS with 0.05% Tween-20 pH 7.4 and then washed 4x, 10 minutes with PBS-Tween 20.
- 5 A mixture of SpA, SpG and SpL at a concentration of 5 µg/ml is added to the membranes.
- 6 After that there is an incubation period of 12 hours at 4°C. It may be an overnight incubation period.
- 7 The nitrocellulose membranes were washed as above.
- 8 A secondary antibody (rabbit anti-chicken IgY horseradish peroxidase, Sigma Aldrich) is added at a 1:15 000 dilution.
- 9 It is incubated for one hour at room temperature and washed as above.
- 10 Tetramethyl-benzidine (TMB) solution is added to the nitrocellulose membranes, which are then incubated in the dark for seven minutes. Then, the membranes are shaken gently and rinsed thoroughly in de-ionized water to stop the blotting process and are left to dry.
- 11 Alternatively, Ig samples are transferred to nitrocellulose membranes and directly probed using SpLAG-HRP (diluted 1:5000) and then adding TMB (this system was mainly used for detecting avian Igs).