



Aug 22, 2022

# Rapid Diagnostic Tests and ELISA for diagnosing Chronic Chagas Disease: Systematic revision and meta-analysis

Sandra Helena Suescún-Carrero<sup>1</sup>, Philippe Tadger<sup>2</sup>, Carolina Sandoval Cuellar<sup>3</sup>, lauramirez<sup>3</sup>, Lluis Armadans-Gil<sup>4</sup>

<sup>1</sup>Universidad de Boyacá, Universitat Autónoma de Barcelona, Barcelona, Spain.;

<sup>2</sup>Real World Solutions, IQVIA, Zaventem, Belgium, Universidad de Boyacá, Tunja, Colombia;

<sup>3</sup>Universidad de Boyacá;

<sup>4</sup>Epidemiology and Preventive Medicine Service, Hospital Universitari Vall d'Hebron - Universitat Autónoma de Barcelona, Barcelona, Spain.



dx.doi.org/10.17504/protocols.io.4r3l2oxppv1y/v1

Sandra Helena Suescún-Carrero



1

ABSTRACT

# Objective

To determine the diagnostic validity of the enzyme-linked immunosorbent assay (ELISA) and Rapid Diagnostic Tests (RDT) among individuals with suspected chronic Chagas Disease (CD).

# Methodology

A search was made for studies with ELISA and RDT assays validity estimates as eligibility criteria, published between May and August 2020 on PubMed, Web of Science, Scopus, and LILACS. This way, we extracted the data and assessed the risk of bias and applicability of the studies using the QUADAS-2 tool. The bivariate random effects model was also used to estimate the overall sensitivity and specificity through *forest-plots*, ROC space, and we visually assessed the heterogeneity between studies. Meta-regressions were made using subgroup analysis. We used Deeks' test to assess the risk of publication bias.

#### Results

43 studies were included; 27 assessed ELISA tests; 14 assessed RDTs; and 2 assessed ELISA and RDTs, against different reference standards. 51.2 % of them used a non-comparative observational design, and 46.5 % a comparative clinical design ("case-control" type). High risk of bias was detected for patient screening and reference standard. The ELISA tests had a sensitivity of 99% (95% CI: 98-99) and a specificity of 98% (95% CI: 97-99); whereas the Rapid Diagnostic Tests (RDT) had values of 95% (95% CI: 94-97) and 97% (95% CI: 96-98), respectively. Deeks' test showed asymmetry on the ELISA assays.

#### **Conclusions**

ELISA and RDT tests have high validity for diagnosing chronic Chagas disease. The analysis of these two types of evidence in this systematic review and meta-analysis constitutes an input for their use. The limitations included the difficulty in extracting data due to the lack of information in the articles, and the comparative clinical-type design of some studies. This article was funded through the Universidad de Boyacá.

**ATTACHMENTS** 

Protocolo RS Chagas.pdf

DOI

dx.doi.org/10.17504/protocols.io.4r3l2oxppv1y/v1



#### PROTOCOL CITATION

Sandra Helena Suescún-Carrero, Philippe Tadger, Carolina Sandoval Cuellar, lauramirez, Lluis Armadans-Gil 2022. Rapid Diagnostic Tests and ELISA for diagnosing Chronic Chagas Disease: Systematic revision and meta-analysis. **protocols.io** 

https://protocols.io/view/rapid-diagnostic-tests-and-elisa-for-diagnosing-ch-cfixtkfn

## FUNDERS ACKNOWLEDGEMENT

Universidad de Boyacá

Grant ID: www.uniboyaca.edu.co

Universitat Autónoma de Barcelona, Barcelona, Spain.

Grant ID: www.UAB.cat

# **KEYWORDS**

Rapid Diagnostic Tests, ELISA, diagnosing Chronic, Chagas Disease, Systematic revision, meta-analysis

## LICENSE

This is an open access protocol distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**CREATED** 

Aug 19, 2022

LAST MODIFIED

Aug 22, 2022

PROTOCOL INTEGER ID

68919

## MATERIALS TEXT

Bases de datos PubMed, Web of Science, Scopus, and LILACS; hojas, esferos, packages (meta4diag: Binomial-normal with probit, and metandi and IGS: Binomial-normal with logit); (R DTAplots program), R 1.3 *software* (DTAplots, bamdit::plotcompare and meta4diag::meta-regression), Stata 15 (metandi), midas and JAGS

# 1 Protocol and registration

# protocols.io

2	Eligibility criteria
3	Data sources
4	Study search and selection
5	Data collection process
6	Definition for data extraction
7	Risk of bias and applicability
8	Diagnostic accuracy measures
9	Summary of results
10	Addinitional analyses
11	