



AUG 24, 2023

# Histology and Retention of Implanted Carbon Fiber Thread Electrodes in Fixed Tissue

Ann M

Helen N Schwerdt<sup>1,2</sup>, Tomoko Yoshida<sup>3</sup>, Graybiel<sup>3</sup><sup>1</sup>Department of Bioengineering, University of Pittsburgh;<sup>2</sup>Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD;<sup>3</sup>McGovern Institute and Department of Brain and Cognitive Sciences, Massachusetts Institute of TechnologyOPEN  ACCESS

Helen N Schwerdt



## ABSTRACT

Methods to cut and retain implanted carbon fiber electrodes in brain tissue are described.

### DOI:

[dx.doi.org/10.17504/protocols.io.6qpvr4nm3gmk/v1](https://dx.doi.org/10.17504/protocols.io.6qpvr4nm3gmk/v1)

**Protocol Citation:** Helen N Schwerdt, Tomoko Yoshida, Ann M Graybiel 2023. Histology and Retention of Implanted Carbon Fiber Thread Electrodes in Fixed Tissue . **protocols.io** <https://dx.doi.org/10.17504/protocols.io.6qpvr4nm3gmk/v1>

**License:** This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working  
We use this protocol and it's working

**Created:** Apr 11, 2023

**Last Modified:** Aug 24, 2023

**PROTOCOL integer ID:**  
80334

- 1 Rats were deeply anesthetized with Euthasol (pentobarbital sodium and phenytoin sodium from Virbac AH Inc.), then transcardially perfused with 0.9% saline solution followed by 4% paraformaldehyde in 0.1 M phosphate buffer (PB).
- 2 The implanted electrodes were cut just above the skull and below the copper wire bundle component of the device with a rotary saw (Dremel) to retain the flexible Py-CF portion of the device in the fixed brain tissue.
- 3 Brains were removed from the skull, post-fixed for 24 hours in the same fixative solution and stored in 25% glycerol with 0.075% sodium azide in 0.1 M PB overnight or until cutting.
- 4 To prepare sections, brains were frozen on dry ice then cut in the transverse plane on a freezing microtome with a microtome knife (C. L. Sturkey, Inc. #K185A).
- 5 Sections were cut sequentially, with four sections at 25- $\mu$ m then one section at 100- $\mu$ m, repeated until the end of the frozen block of brain.
- 6 The 100- $\mu$ m sections were immediately mounted onto glass slides and air dried, whereas the 25- $\mu$ m sections were stored in 0.1% sodium azide in 0.1 M PB at 4°C until use.