



May 06, 2020

Protocol S-100B Determination in Melanoma

Samantha Damude¹, Schelto Kruijff¹, Harald J Hoekstra¹, Kevin P Wevers¹¹University Medical Center Groningen

1 Works for me dx.doi.org/10.17504/protocols.io.bf4cqsw

Samantha Damude

ABSTRACT

This protocol describes the implementation of a dummytube system in AJCC stage III melanoma patients, to investigate whether serum S-100B levels can be falsely elevated by adipocyte contamination in the first drawn serum separation tube.

A dummy tube system was introduced for performance of venipunctures during regular follow-up in all AJCC stage III melanoma patients, with no evidence of disease. After venipuncture, a dummy tube was drawn first, and subsequently the regular tube was drawn. The first tube was anonymously coded and stored, while the second tube was registered in the patients' medical results. After performing the assay, S-100B levels between the two consecutively drawn samples were compared.

THIS PROTOCOL ACCOMPANIES THE FOLLOWING PUBLICATION

Double Venipuncture is not Required for Adequate S-100B Determination in Melanoma Patients.

GUIDELINES

Background

Patients with AJCC stage I / II melanoma have a 20-30% chance of a positive sentinel node.^{1,2} Regional gland metastases develop in 20-60% of patients with stage I / II melanoma^{1,3} and 10-50% of these patients eventually develop distant metastases.^{3,4,5}

Patients with clinical stage III melanoma are staged using an FDG-PET and / or CT scan. Distant metastases appear to be present in 32% of these patients with palpable lymph nodes and are therefore stage IV.^{6,7} The surgical treatment of regional gland metastases is a complementary or therapeutic lymph node dissection. The surgical techniques for the various gland dissections have been defined.⁸ The indication for additional radiotherapy is also defined.⁹

Research in stage III melanoma has shown that the biomarker S-100B has a predictive value with respect to disease-free and melanoma specific survival; at normal values ($<0.20\mu\text{g} / \text{l}$) 38% disease-free and 47% 5-year survival, at elevated values 7% disease-free and 28% 5-year survival, respectively.^{5,10} Early detection of distant metastases and, if possible, surgical treatment of these, contributes to an improved 5-year survival (15-28% 5-year survival with surgical treatment, as opposed to 5-10% with system therapy, based on data from the years 1993 to 2008).¹¹

The biomarker S-100B appears to be more reliable than LDH in the early detection of melanoma metastases.¹⁰ In recent years, there has been a real breakthrough in the treatment of metastatic melanoma in the field of "targeted therapy" and in the field of "immunotherapy".¹² Early diagnosis of melanoma metastases is generally associated with lower tumor load and therefore potentially more successful treatment.¹³

National guideline melanoma follow-up

The Dutch Melanoma Guideline does not, except for the frequency of follow-up, contain any advice regarding additional diagnostics to be carried out during the follow-up, in contrast to other National guidelines.^{14,15} As outlined earlier, a biomarker is available, S-100B, which can indicate the presence or absence of metastatic

disease.^{5,16} This biomarker has been implemented in the German Melanoma Guideline, due to a higher sensitivity for disease progression in melanoma than LDH.¹⁷

In anticipation of a possible future change of the Dutch Melanoma Guideline, the Department of Surgical Oncology at the University Medical Center Groningen has decided to amend the Protocol Follow-up Melanoma stage III, given the successful breakthroughs that have been achieved in the systemic treatment of the melanoma. These treatments are available for recognized melanoma centers in the Netherlands, including the UMCG.

Protocol Follow-up Melanoma Stage III UMCG

	Clinical visit + LDH + S-100B
1st year	4 times per year
2nd year	3 times per year
3rd – 5th year	2 times per year
> 5th year	1 times per year

The laboratory results of S-100B and LDH are registered in the Electronic Patient Record. The reference value of S-100B is <0.20µg / l and of LDH <250U / l.

At a marginally increased value of S-100B (0.20-0.30µg / l) and / or LDH (250-400U / l), the S-100B and LDH will be reassessed after 4 weeks. At an absolutely increased value of S-100B (> 0.30 µg / l) and / or LDH (> 400U / l) an FDG-PET / CT will be performed and an MRI scan of the brain in case of FDG positive hearths. In addition, standard mutation analyzes will be requested in stage III melanoma patients: BRAF, MET, C KIT. Patients with elevated biomarkers are discussed in the multidisciplinary melanoma discussion in accordance with the SONCOS guideline.¹⁸

BEFORE START

The study was conducted in accordance with the Declaration of Helsinki, and conforms to the guidelines of the central medical ethics committee (**METc2015.215**).

MATERIALS TEXT

Determination of S-100B in a second blood tube (dummytube)

Recent research by the UMCG has shown that drawing a second tube of blood before the S-100B tube can prevent possible incorrect values.¹⁹ For the careful introduction of this extra serum tube, we ask the patients' permission to determine the S-100B value in this tube as well. This extra result will be processed anonymously and will not be disclosed to the patient. By handing over the extra tube to the puncture lab, the patient agrees to the additional determination.

If patients do not want to participate, they can return the tube to the outpatient clinic.

Patient ID number	
Puncture date	
Project number	
S-100B result project	
Routine number	
S-100B result routine	
>1 cm subcutaneous	Yes / No

Traumatic puncture	Yes / No
--------------------	----------

S-100B analysis

S-100B concentrations were determined by performing the S-100B assay (Diasorin, Saluggia, Italy) on an ELISA Robot platform (DS2, Dynex Technologies, Magellan Biosciences, Worthing, United Kingdom). After routine centrifugation, serum was separated from the tubes, aliquoted and stored at -80 °C. The two tubes of all individuals were analyzed after thawing the samples, using the S-100B ELISA according to manufacturer's protocol.

The intra-assay Coefficient of Variation of the S-100B assay is 7% at levels of 0.04 µg/L (0.0028 µg/L). The Limit of Blank was determined to be 0.0034 µg/L, whereas the Limit of Quantitation (20 % CV) was determined to be 0.092 µg/L.

The reference interval was determined by analysis of S-100B values in 120 healthy individuals (median 0.07; range 0.01-0.59) and calculating the 95% confidence interval according to the Clinical and Laboratory Standards Institute EP28-A3c guideline (formerly C28-A2), resulting in a reference cut-off value for the healthy population of 0.20 µg/L.²⁰

References

1. Morton DL. Overview and update of the phase III multicenter selective lymphadenectomy trials (MSLT-I and MSLT-II) in melanoma. Clin Exp Metastasis 2012 Oct;29(7):699-706.
2. Speijers MJ, Bastiaannet E, Sloot S, Suurmeijer AJ, Hoekstra HJ. Tumor mitotic rate added to the equation: Melanoma prognostic factors changed? : A single-institution database study on the prognostic value of tumor mitotic rate for sentinel lymph node status and survival of cutaneous melanoma patients. Ann Surg Oncol 2015 Jan 21.
3. Francken AB, Bastiaannet E, Hoekstra HJ. Follow-up in patients with localised primary cutaneous melanoma. Lancet Oncol 2005 Aug;6(8):608-21.
4. Francken AB, Accortt NA, Shaw HM, Wiener M, Soong SJ, Hoekstra HJ, Thompson JF. Prognosis and determinants of outcome following locoregional or distant recurrence in patients with cutaneous melanoma. Ann Surg Oncol 2008 May;15(5):1476-84.
5. Kruijff S, Hoekstra HJ. The current status of S-100B as a biomarker in melanoma. Eur J Surg Oncol 2012 Apr;38(4):281-5.
6. Bastiaannet E, Wobbes T, Hoekstra OS, van der Jagt EJ, Brouwers AH, Koelemij R, de Klerk JM, Oyen WJ, Meijer S, Hoekstra HJ. Prospective comparison of [18F]fluorodeoxyglucose positron emission tomography and computed tomography in patients with melanoma with palpable lymph node metastases: Diagnostic accuracy and impact on treatment. J Clin Oncol 2009 Oct 1;27(28):4774-80.
7. Niebling MG, Bastiaannet E, Hoekstra OS, Bonenkamp JJ, Koelemij R, Hoekstra HJ. Outcome of clinical stage III melanoma patients with FDG-PET and whole-body CT added to the diagnostic workup. Ann Surg Oncol 2013 Sep;20(9):3098-105.
8. Balch CM, Houghton AN, Sober AJ, Soong SJ, Atkin MB, Thompson JF. Cutaneous melanoma, fifth edition. St. Louis, MO: Quality Medical Publishing, Inc.; 2009. .
9. Burmeister BH, Henderson MA, Ainslie J, Fisher R, Di Iulio J, Smithers BM, Hong A, Shannon K, Scolyer RA, Carruthers S, et al. Adjuvant radiotherapy versus observation alone for patients at risk of lymph-node field relapse after therapeutic lymphadenectomy for melanoma: A randomised trial. Lancet Oncol 2012 Jun;13(6):589-97.
10. Wevers KP, Kruijff S, Speijers MJ, Bastiaannet E, Muller Kobold AC, Hoekstra HJ. S-100B: A stronger prognostic biomarker than LDH in stage IIIB-C melanoma. Ann Surg Oncol 2013 Aug;20(8):2772-9.

11. Wevers KP, Hoekstra HJ. Stage IV melanoma: Completely resectable patients are scarce. *Ann Surg Oncol* 2013 Jul;20(7):2352-6.
12. Stadler S, Weina K, Gebhardt C, Utikal J. New therapeutic options for advanced non-resectable malignant melanoma. *Adv Med Sci* 2014 Dec 25;60(1):83-8.
13. Leiter U, Buettner PG, Eigentler TK, Forschner A, Meier F, Garbe C. Is detection of melanoma metastasis during surveillance in an early phase of development associated with a survival benefit? *Melanoma Res* 2010 Jun;20(3):240-6.
14. Melanoom, Landelijke Richtlijn, Versie: 2.0 [Internet]; c201213-08-2012]. Available from: <http://www.oncoline.nl/melanoom>
15. Speijers MJ, Francken AB, Hoekstra-Weebers JEHM, Bastiaannet E, Kruijff S, Hoekstra HJ. Optimal follow-up for melanoma. *Expert Rev Dermatol* 2010 ;5(4):461-78.
16. Beyeler M, Waldispuhl S, Strobel K, Joller-Jemelka HJ, Burg G, Dummer R. Detection of melanoma relapse: First comparative analysis on imaging techniques versus S100 protein. *Dermatology* 2006;213(3):187-91.
17. Dummer R, Hauschild A, Guggenheim M, Keilholz U, Pentheroudakis G, ESMO Guidelines Working Group. Cutaneous melanoma: ESMO clinical practice guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2012 Oct;23 Suppl 7:vii86-91.
18. SONCOS; Multidisciplinaire normering oncologische zorg in Nederland [Internet]; Februari 2014. Available from: <https://www.iknl.nl/docs/default-source/Palliatieve-zorg-in-de-ziekenhuizen/soncos.pdf?sfvrsn=0>
19. Damude S, Niebling MG, Kobold AC, Hoekstra HJ, Kruijff S, Wevers KP. Adipocytes in venipunctures cause falsely elevated S-100B serum values. *Clin Chem Lab Med* 2016;54:e235-7.
20. Horowitz GL. CLSI EP28-A3c: Defining, Establishing, and Verifying Reference Intervals in the Clinical Laboratory; Approved Guideline – Third Edition. Available at: <http://shop.clsi.org/s.nl/it.A/id.257/f?sc=7&category=2395&sc=7&category=2395>