

6



Apr 18, 2022

© Depletion of mtDNA with Ethidium Bromide

William Hancock-Cerutti^{1,2,3,4,5}, Pietro De Camilli^{1,2,3,5}

¹Departments of Neuroscience and of Cell Biology, Yale University School of Medicine, New Haven, Connecticut 06510, USA;

²Howard Hughes Medical Institute;

³Program in Cellular Neuroscience, Neurodegeneration and Repair, Yale University School of Medici ne, New Haven, Connecticut 06510, USA;

⁴Interdisciplinary Neuroscience Program and MD-PhD Program, Yale University School of Medicine, New Haven, Connecticut 06510, USA;

⁵Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD, 20 815



dx.doi.org/10.17504/protocols.io.rm7vzy6e5lx1/v1

William Hancock-Cerutti

This protocol describes the depletion of mitochondrial DNA (mtDNA) from HeLa cells using ethidium bromide (EtBr).

HC-Protocol-mtDNAdepletion.docx

DOI

dx.doi.org/10.17504/protocols.io.rm7vzy6e5lx1/v1

William Hancock-Cerutti, Pietro De Camilli 2022. Depletion of mtDNA with Ethidium Bromide. **protocols.io**

https://dx.doi.org/10.17504/protocols.io.rm7vzy6e5lx1/v1

_____ protocol,

Apr 18, 2022

Apr 18, 2022

60898



1