




MAY 23, 2023

# Collecting supernatant from C. elegans culture and lyophilizing supernatant

Man  
Muhammad Zaka Asif<sup>1</sup>, Shah<sup>1</sup>, Yosef Smadi<sup>1</sup>

<sup>1</sup>Edison Lab, UGA



Muhammad Zaka Asif

OPEN ACCESS

**DOI:**  
[dx.doi.org/10.17504/protocols.io.36wgqjxxvk5/v1](https://dx.doi.org/10.17504/protocols.io.36wgqjxxvk5/v1)

**Protocol Citation:** Muhammad Zaka Asif, Man Shah, Yosef Smadi 2023. Collecting supernatant from C. elegans culture and lyophilizing supernatant. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.36wgqjxxvk5/v1>

**License:** This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working  
We use this protocol and it's working

**Created:** May 23, 2023

**Last Modified:** May 23, 2023

**PROTOCOL integer ID:**  
82342

## ABSTRACT

This protocol describes day 5 of our workflow to grow worms in liquid culture to induce the production of natural products (in this case, 1-HP derivatives). In this protocol, we get an approximate count for the number of life-stage synchronized worms exposed to the toxin, and we collect the supernatant, which contains 1-HP and its derived metabolites. After collecting the supernatant, we freeze and lyophilize it to prepare for extraction.

- 1 Remove the flasks from the shaker and estimate their population by aliquoting onto glass slides as before.

- 2** Centrifuge at 525 RCF for 1 minute and collect the supernatant in multiple 2 mL microcentrifuge tubes.
- 3** Save the worm pellet at -80°C if desired for future studies.
- 4** Centrifuge the microcentrifuge tubes at 20,800 RCF for 10 minutes. This will separate the bacteria from the rest of the supernatant (bacteria will settle to the bottom).
- 5** Combine all the supernatant remaining from the microcentrifuge tubes once the spin is completed in one 50 mL conical tube.
- 6** Freeze the supernatant with liquid nitrogen and lyophilize for 24-40 hrs until complete.