



Aug 23, 2020

Copy of Copy of Copy of ELISA for quantification of human C9 in serum or plasma.

Angel A Justiz-Vaillant¹, Belkis Ferrer-Cosme²

¹University of the West Indies St. Augustine; ²Saturnino Lora Torres' Provincial Teaching Clinical Surgical Hospital. Cuba

In Development

dx.doi.org/10.17504/protocols.io.bj78krw



Angel Justiz-Vaillant
University of the West Indies St. Augustine

DOI

dx.doi.org/10.17504/protocols.io.bj78krw

PROTOCOL CITATION

Angel A Justiz-Vaillant, Belkis Ferrer-Cosme 2020. Copy of Copy of Copy of ELISA for quantification of human C9 in serum or plasma.. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bj78krw>

LICENSE

This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Aug 23, 2020

LAST MODIFIED

Aug 23, 2020

PROTOCOL INTEGER ID

40928

- 1 An anti-human C9 coating antibody is adsorbed onto the microwells by incubation overnight at 4°C with carbonate-bicarbonate buffer.
- 2 Add 50 µl of human serum or plasma. Human C9 present in the serum or plasma binds to antibodies adsorbed into the microwells.
- 3 The microplate is blocked with 3% non-fat milk-PBS buffer and later wash to remove unbound proteins.
- 4 Fifty (50) µl of biotin-conjugated anti-C9 antibody is added. The optimal dilution must be investigated.
- 5 The microplate is rewashed with PBS-Tween 20 buffer, pH 7.4.

- 6 One hundred μ l of streptavidin-HRP conjugate is added and it binds to the biotin-conjugated anti-C9 antibody. The optimal dilution of this conjugate must be investigated.
- 7 The plate is washed following incubation to remove the unbound Streptavidin-HRP.
- 8 Add 100 μ l of 3,3',5,5'- tetramethylbenzidine (TMB; Sigma-Aldrich) into each well.
- 9 Incubate the microwells in the dark for 15 min.
- 10 A colored product is formed in proportion to the quantity of C9 present in the sample or standard.
- 11 The reaction is terminated by addition of 100 μ l 3M H₂SO₄ and the absorbance is measured at 450 nm.
- 12 A standard curve is made from 7 human C9 standard dilutions and the human C9 sample concentration is determined.