

May 28, 2024

Accelerated Ovarian Failure AOF mouse model: a model of transitional menopause

DOI

dx.doi.org/10.17504/protocols.io.yxmvmnkbng3p/v1

Roberta Marongiu¹

¹Weill Cornell Medical College

ASAP Collaborative Rese...

Kaplift Protocols



Eileen Ruth Torres

Weill Cornell Medicine

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.yxmvmnkbng3p/v1

Protocol Citation: Roberta Marongiu 2024. Accelerated Ovarian Failure AOF mouse model: a model of transitional menopause. protocols.io <https://dx.doi.org/10.17504/protocols.io.yxmvmnkbng3p/v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: January 25, 2022

Last Modified: May 28, 2024

Protocol Integer ID: 57412

Keywords: ASAPCRN

Funders Acknowledgement:

Aligning Science Across

Parkinson's

Grant ID: 020608

Abstract

Until recently, aging and ovariectomy (OVX) have been the primary rodent models available to examine effects of ovarian hormone loss. These models fail to adequately recapitulate the hormonal changes of the human menopause process. The intact aging model fails to achieve very low estrogen levels at post-menopause, and the OVX model lacks a peri-menopause phase. Conversely, the novel model of AOF successfully replicates the human peri- and post-menopause stages, including erratic estrous cycles and hormone fluctuations, followed by undetectable estrogen levels[1, 2]. In the AOF model, the ovotoxin 4-vinylcyclohexene diepoxide (VCD, Sigma-Aldrich) is injected (130 mg/kg in 0.5% dimethyl sulfoxide in sesame oil; 5 days/week for 3 weeks; i.p.) in mice which induces selective depletion of ovarian primary and primordial follicles[3]. Previous studies from the Milner lab and other groups confirmed that VCD administration does not negatively affect peripheral tissues nor crosses the blood-brain barrier [1, 2]. In addition to reducing confounds associated with surgical manipulations, the AOF model maintains the presence of ovarian tissue which importantly parallels to human menopause. Based on assessment of ovarian follicle depletion and responses of plasticity markers in brain areas known to be estrogen responsive, timepoints for pre-, peri- and post-AOF have been established [1-4]. Timepoints in the AOF model include peri-AOF (>58 days) and post-AOF (>139 days) corresponding to human peri- and post-menopause respectively [1-5].

References

1. Van Kempen, T.A., et al., *Characterization of neural estrogen signaling and neurotrophic changes in the accelerated ovarian failure mouse model of menopause*. Endocrinology, 2014. **155**(9): p. 3610-23.
2. Van Kempen, T.A., T.A. Milner, and E.M. Waters, *Accelerated ovarian failure: a novel, chemically induced animal model of menopause*. Brain Res, 2011. **1379**: p. 176-87.
3. Lohff, J.C., et al., *Effect of duration of dosing on onset of ovarian failure in a chemical-induced mouse model of perimenopause*. Menopause, 2006. **13**(3): p. 482-8.
4. Mayer, L.P., et al., *The follicle-deplete mouse ovary produces androgen*. Biol Reprod, 2004. **71**(1): p. 130-8.
5. Marques-Lopes, J., et al., *Redistribution of NMDA Receptors in Estrogen-Receptor-beta-Containing Paraventricular Hypothalamic Neurons following Slow-Pressor Angiotensin II Hypertension in Female Mice with Accelerated Ovarian Failure*. Neuroendocrinology, 2017. **104**(3): p. 239-256.

Materials

Reagents:

DMSO



sesame oil

 ovotoxin 4-vinylcyclohexene diepoxide (VDC) **Sigma Aldrich Catalog #94956**




Ovotoxin 4-vinylcyclohexene diepoxide (VDC) preparation

2h 30m

- 1 Determine the number of mice to treat with VCD to calculate how much VCD is needed each day.
- 2 Weigh all the mice and obtain an average weight at least once a week for the duration of the VCD treatment. This works only if mice have very similar weights (+/- a few grams around the average). It is recommended to use the high end of variation for calculations to make sure the heaviest mice get enough VCD.
- 3 In the chemical hood, measure volume of VCD from new vial.
- 4 Calculate how much sesame oil to add to the VCD vial in order to prepare the working solution (130 mg/kg).
- 5 Add sesame oil using a pipet with wide opening trying to avoid bubbles which can affect your volume measurements.
- 6 Stir solution for  02:00:00 . 2h
- 7 Parafilm well and store it in the fridge.
- 8 Every day before use, stir vial for  00:30:00 again. 30m

VDC injections



5 days / week for 3 weeks = i.p injection ; 130 mg/kg in 0.5% dimethyl sulfoxide in sesame oil

