




NOV 28, 2023

Transformation of Plasmid into Competent E. coli Cells (e.g., DH5α)

Hussain Zubair¹

¹ZZU



Hussain Zubair

ABSTRACT

This is a detailed experimental protocol for the transformation of plasmid DNA into competent E. coli cells, specifically using the DH5α strain. The protocol encompasses the entire process, from the preparation of LB agar plates supplemented with ampicillin to the final inoculation of transformed cells in LB broth. Key steps include the careful thawing of competent cells, the addition of plasmid DNA, a precise heat shock treatment at 42°C for enhanced plasmid uptake, and subsequent recovery and plating of the cells. The methodology is designed to maximize the efficiency of transformation, crucial for subsequent molecular biology experiments such as gene cloning and expression analysis. This protocol serves as a fundamental guide for researchers in the field of molecular genetics and microbiology.

OPEN ACCESS



DOI:

dx.doi.org/10.17504/protocols.io.eq2lyj2ywlx9/v1

Protocol Citation: Hussain Zubair 2023. Transformation of Plasmid into Competent E. coli Cells (e.g., DH5α).

protocols.io

<https://dx.doi.org/10.17504/protocols.io.eq2lyj2ywlx9/v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Nov 28, 2023

Preparation of LB Agar Plates with Antibiotics

- Prepare LB (Luria-Bertani) agar plates.
 - Once the agar has cooled but not solidified, add ampicillin at a ratio of 1:1000 to the agar.
 - Pour the LB agar into 90 mm Petri dishes and allow to solidify.

Thawing of Competent Cells

10m

- Retrieve the competent E. coli cells (e.g., DH5α strain) from the -80 °C freezer.
 - Thaw the cells slowly on ice to ensure they remain in liquid form. 00:10:00

Addition of Plasmid

30m

- Add 1 µL of the plasmid solution to the competent cells. The volume may vary depending on plasmid's concentration.
 - Incubate the mixture on ice for 00:30:00 min to allow the cells to fully absorb the plasmid.

Heat Shock Treatment

1m 10s

- Subject the cell-plasmid mixture to heat shock in a water bath set at 42 °C for exactly 00:01:10 min.
 - This step facilitates the uptake of the plasmid by the cells.





Cooling Period (Stabilizing the Cells)

- Immediately after the heat shock, place the mixture on ice for 00:02:00 min to stabilize the cells. 2m

Plating on LB Agar

18h

- 6
 - Gently spread the competent cell-plasmid mixture onto the prepared LB agar plates with ampicillin
 - Incubate the plates at  37 °C  Overnight to allow for bacterial growth and colony formation.

18h

Further Inoculation

- 7
 - The following day, select a single colony from the agar plate.
 - Inoculate the colony in LB broth for further growth and experimentation as required