

May 03, 2024

Transmission Electron Microscopy of Native Nanodiscs

DOI

dx.doi.org/10.17504/protocols.io.x54v92ydml3e/v1

Caroline Brown^{1,2,3}, Snehasish Ghosh^{1,2,3}, Kallol Gupta^{1,2,3}

- ¹Nanobiology Institute, Yale University, West Haven, CT, USA;
- ²Department of Cell Biology, Yale University School of Medicine, New Haven, CT, USA;
- ³Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD, 20815



Caroline Brown

Yale University

OPEN ACCESS



DOI: dx.doi.org/10.17504/protocols.io.x54v92ydml3e/v1

Protocol Citation: Caroline Brown, Snehasish Ghosh, Kallol Gupta 2024. Transmission Electron Microscopy of Native Nanodiscs. protocols.io https://dx.doi.org/10.17504/protocols.io.x54v92ydml3e/v1

License: This is an open access protocol distributed under the terms of the <u>Creative Commons Attribution License</u>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's

working

Created: May 03, 2024

Last Modified: May 03, 2024

Protocol Integer ID: 99194

Abstract

This is a protocol for conducting transmission electron microscopy of native nanodiscs to determine population size distribution and morphology.



Staining grids

32m 35s

- Dilute nanodisc samples into single molecule range, typically around between 1:10 and 1:100 is the samples have been through size exclusion chromatography using dilution buffer (

 [M] 50 millimolar (mM) Tris HCl pH 7.4)
- 2 Glow-discharge carbon-coated copper grids (200 mesh) for 00:00:30 seconds.

30s

1m

4 Wash grid once with Δ 5 μL uranyl formate for 🕥 00:00:05 seconds.

5s

Stain grid with Δ 5 µL uranyl formate for 00:01:00 minute and blot dry with Whiteman ashless filter paper.

1m

6 Allow to dry for 00:30:00 minutes to Overnight

30m

Image Acquisition

Take micrographs using a JEOL JEM 1400PLUS electron microscope at an operating voltage of 80 kV.