

OCT 04, 2023

# OPEN ACCESS



DOI:

dx.doi.org/10.17504/protocol s.io.8epv5xe9jg1b/v1

**Protocol Citation:** John Sembrat, Marta Bueno, Alison Morris, koenigshoffm, Oliver Eickelberg 2023. UPitt TriState SenNet TMC Heart Procurement. **protocols.io** https://dx.doi.org/10.17504/p rotocols.io.8epv5xe9jg1b/v1

License: This is an open access protocol distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working We use this protocol and it's working

Created: Oct 04, 2023

Last Modified: Oct 04, 2023

**PROTOCOL** integer ID:

88805

Keywords: heart

### ( ) UPitt TriState SenNet TMC Heart Procurement

John Oliver Sembrat<sup>1</sup>, Marta Bueno<sup>1</sup>, Alison Morris<sup>1</sup>, koenigshoffm<sup>1</sup>, Eickelberg<sup>1</sup>

<sup>1</sup>Division of Pulmonary and Critical Care Medicine. Department of Medicine. School of Medicine

TriState SenNet

Cellular Senescence Network (SenNet) Method Development Community



Marta Bueno

Division of Pulmonary and Critical Care Medicine. Department...

#### **ABSTRACT**

This document outlines the request for specimen procurement before arrival at the TriState SenNet TMC Biospecimen Core at the University of Pittsburgh, as part of the Cellular Senescence Network Program (SenNet).

### Requested processing protocol for Donation following Brain Death Decla...

- Organ Procurement Organization Recovery Team prepares for organ recovery as consistent with organ transplant protocol with cross clamp and in situ cold flush body cooling with minimal warm ischemic time.
- 2 Heart retrieval: The superior caval vein is ligated. The inferior caval vein is opened and the left atrium is vented, in order to prevent left ventricular distension. The ascending aorta is subsequently cross clamped and cold cardioplegia is administered via the aortic root to achieve hypothermic diastolic arrest.
- 3 Place Heart in a Specimen Bag and submerge with chilled Perfadex or similar. Triple bag the tissue. (if recover with lungs)

if retrieved alone, heart is packed in one or two litres of cardioplegia and placed in an ice box for transport.

## Requested processing protocol for Donation following Cardiac Death /Ar...

- 4 Organ Procurement Organization / Recovery Team prepares for organ recovery as quickly following Cardiac Arrest while in comfort care as authorized by next of kin.
- Remainder of organ / tissue recovery should be as consistent with organ transplant protocol (section above) as possible.

Includes organ flush with Perfadex or similar organ transplant buffer until solution runs clear.

6 All warm ischemic time is to be minimized and reported. Tissues may be accepted up to 6 hours WIT. Other WIT may be evaluated depending of the circumstances.