

Jul 11, 2024

Assessing Astrocyte Territory Volume and 3D Sholl Analysis

DOI

dx.doi.org/10.17504/protocols.io.rm7vzj734lx1/v1

Shiyi Wang¹

¹Duke University

ASAP Collaborative Rese...



Shiyi Wang

Duke University

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.rm7vzj734lx1/v1

Protocol Citation: Shiyi Wang 2024. Assessing Astrocyte Territory Volume and 3D Sholl Analysis. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.rm7vzj734lx1/v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: July 11, 2024

Last Modified: July 11, 2024

Protocol Integer ID: 103198

Keywords: ASAPCRN

Funders Acknowledgement:
Aligning Science Across
Parkinson's (ASAP) initiative
Grant ID: ASAP-020607



Disclaimer

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to **protocols.io** is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with **protocols.io**, can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

Abstract

Assessing Astrocyte Territory Volume and 3D Sholl Analysis



1 ****Section Collection****

- 1.1 - Collect 100 μm -thick floating sections containing the anterior cingulate cortex (ACC) and primary motor cortex (MOp) from mice.
- 1.2 - Ensure astrocytes are sparsely labeled via PALE with mCherry-CAAX.

2 ****Imaging****

- 2.1 - Use an Olympus FV 3000 microscope with a 60x objective to acquire high-magnification images (50-60 μm z-stack) of individual astrocytes.
- 2.2 - Criteria for inclusion: Capture the entire astrocyte within a single brain section, located in layer 2/3 of ACC or MOp.

3 ****Surface Reconstructions****

- 3.1 - Analyze imaged astrocytes using Imaris Bitplane software.
- 3.2 - Generate surface reconstructions of the astrocytes.

4 ****Territory Volume Measurement****

- 4.1 - Use Imaris Xtensions “Visualize Surface Spots” and “Convex Hull” to create additional surface renders representing the territory of each astrocyte.
- 4.2 - Record the volume of each astrocyte territory.

5 ****Statistical Testing****



- 5.1 - Analyze astrocyte territory sizes from biological replicates across experimental conditions.
- 5.2 - Use a nested two-way ANOVA followed by the Bonferroni posthoc test for statistical comparisons.
- 6 ****Surface and Filament Creation****
- 6.1 - Load astrocyte images onto Imaris and create a surface representation for each astrocyte.
- 7 ****Filament Generation****
- 7.1 - Create filaments using 'Add new filament (leaf icon)' in Imaris.
- 8 ****Quantification of Complexity****
- 8.1 - Use the gear tool on Imaris to display Sholl intersections for each astrocyte.
- 8.2 - Quantify the complexity of astrocytes based on Sholl intersections.
- 9 ##### Experimental Details
- 10 ### Notes:
- 11 - Maintain consistency in imaging and analysis methodologies across experimental conditions.
- 12 - Consider biological variability and replicate experiments to ensure robust conclusions.