

Jul 10, 2024

Rat astrocyte isolation and culture

DOI

dx.doi.org/10.17504/protocols.io.n2bvjnpxpqk5/v1

Shiyi Wang¹

¹Duke University

ASAP Collaborative Rese...



Shiyi Wang

Duke University

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.n2bvjnpxpqk5/v1

Protocol Citation: Shiyi Wang 2024. Rat astrocyte isolation and culture. **protocols.io**

<https://dx.doi.org/10.17504/protocols.io.n2bvjnpxpqk5/v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: July 10, 2024

Last Modified: July 10, 2024

Protocol Integer ID: 103187

Keywords: ASAPCRN

Funders Acknowledgement:
Aligning Science Across
Parkinson's (ASAP) initiative
Grant ID: ASAP-020607



Disclaimer

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to **protocols.io** is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with **protocols.io**, can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

Abstract

Rat astrocyte isolation and culture

- 1 ****Cortex Dissection and Digestion**** - Micro-dissect P1 rat cortices from both sexes. - Digest cortices in papain solution.
- 2 ****Trituration and Resuspension**** - Triturate tissue in low and high ovomucoid solutions. - Resuspend cells in astrocyte growth media (AGM): - DMEM (GIBCO 11960) - 10% FBS - 10 μ M hydrocortisone - 100 U/ml Pen/Strep - 2 mM L-Glutamine - 5 μ g/ml Insulin - 1 mM Na Pyruvate - 5 μ g/ml N-Acetyl-L-cysteine
- 3 ****Cell Plating and Incubation**** - Plate 15-20 million cells on 75 mm² flasks (non-ventilated cap) coated with poly-D-lysine. - Incubate at 37°C in 10% CO₂.
- 4 ****Non-Astrocyte Cell Removal (DIV 3)**** - Forcefully shake closed flasks by hand for 10-15 seconds to remove non-astrocyte cells. - Ensure only an adherent monolayer of astrocytes remains.
- 5 ****AraC Treatment (DIV 5 to DIV 7)**** - Add AraC to the media to eliminate contaminating fibroblasts.
- 6 ****Astrocyte Passage and Plating (DIV 7)**** - Trypsinize astrocytes with 0.05% Trypsin-EDTA. - Plate cells into 12-well or 6-well dishes.
- 7 ****Transfection of Astrocytes (DIV 8)**** - Transfect with shRNA and/or expression plasmids using Lipofectamine LTX with Plus Reagent (Thermo Scientific). - Dilute 1 μ g (12-well) or 2 μ g (6-well) total DNA in Opti-MEM with Plus Reagent. - Mix with Opti-MEM containing LTX (1:2 DNA to LTX ratio). - Incubate for 30 minutes at room temperature. - Add transfection solution to astrocyte cultures and incubate at 37°C for 3 hours.
- 8 ****Co-Culture with Neurons (DIV 10)**** - Trypsinize astrocytes on DIV 10. - Resuspend in NGM Plus. - Plate 20,000 cells per well onto DIV 10 neurons. - Co-culture for 48 hours.