



OCT 07, 2023

General Guidelines for Culture of Multiple Myeloma Cell Lines

Dennis Juarez¹, dfruman¹

¹University of California, Irvine

Fruman Lab



Dennis Juarez

OPEN ACCESS



DOI:
dx.doi.org/10.17504/protocols.io.rm7vzb6n4vx1/v1

Protocol Citation: Dennis Juarez, dfruman 2023. General Guidelines for Culture of Multiple Myeloma Cell Lines. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.rm7vzb6n4vx1/v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Dec 21, 2022

DISCLAIMER

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to [protocols.io](#) is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with [protocols.io](#), can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

ABSTRACT

General guidelines for culture and use of myeloma cell lines in the Fruman Lab.

MATERIALS

RPMI 1640 (Hyclone SH30255)
 Pen/Strep/Glutamine (Gibco 10378016)
 1M HEPES (Corning 25-060-CI)

- 1 Incubator Settings:** 37 degrees Celsius, 5% CO₂

- 2 Medium:** 88% RPMI 1640, 10% FBS, 1% Pen/Strep/Glutamine, 1% (Final: 10 mM) of 1M HEPES

- 3 Cryopreservation:** Freeze medium: 90% FBS, 10% DMSO. 5 million cells are resuspended in 500uL of freeze media, transferred to cryotubes, and frozen slowly in Mr. Frosty™ Freezing Container at -80 degrees Celsius.

- 4 Guidelines for Experimental Use:**
 1. Passages are not intuitive for suspension cells and passage number should instead be used to indicate number of times the original stock has been grown out and frozen for future experiments.
 2. New cell lines should be grown to at least 35 million cells and a minimum of 6 vials should be frozen for experimental use throughout the year and keeping at least one for long term storage in liquid nitrogen.
 3. Cell lines should be authenticated through STR profiling annually or whenever a new set of stocks are cryopreserved for future experiments to confirm lack of cross contamination.
 4. Cell lines should not be cultured for longer than 2 months when used for experiments.
 5. Cell lines should be maintained at concentrations between 0.5 million to 2 million per mL throughout culture as to not exhaust media.
 6. Experiments requiring incubation times less than 24 hours should have cells plated at 1 million per mL media.
 7. Experiments requiring incubation times greater than 24 hours should have cells plated at 0.25-0.5 million per mL media.
 8. When ordering new lots of FBS, a sample should be requested to test if the new lot will be deleterious to cell survival and cell proliferation.