

May 09, 2024

Cathepsin D assay to verify the retention of lysosomal content

DOI

dx.doi.org/10.17504/protocols.io.8epv5r7j5g1b/v1

Rotimi Y. Fasimoye^{1,2}, Dario R. Alessi^{1,2}

¹Aligning Science Across Parkinson's Collaborative Research Network, Chevy Chase, MD 20815;

²Medical Research Council Protein Phosphorylation and Ubiquitylation Unit, School of Life Sciences, University of Dundee, Dow Street, Dundee DD1 5EH, UK

ASAP Collaborative Rese...



Francesca Tonelli

MRC-PPU at The University of Dundee

OPEN  ACCESS



DOI: **dx.doi.org/10.17504/protocols.io.8epv5r7j5g1b/v1**

Protocol Citation: Rotimi Y. Fasimoye, Dario R. Alessi 2024. Cathepsin D assay to verify the retention of lysosomal content. protocols.io **<https://dx.doi.org/10.17504/protocols.io.8epv5r7j5g1b/v1>**

License: This is an open access protocol distributed under the terms of the **[Creative Commons Attribution License](#)**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: March 14, 2024

Last Modified: May 09, 2024

Protocol Integer ID: 96905

Keywords: ASAPCRN

Funders Acknowledgement:
Aligning Science Across
Parkinson's
Grant ID: ASAP-000463

Disclaimer

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to **protocols.io** is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with **protocols.io**, can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

Abstract

Cathepsin D assay is a fluorescence-based assay that leverage on the activity of cathepsin D, a lysosomal enzyme, to monitor the intactness of lysosome in the cell. Here, we describe a method where we used the measurement of cathepsin D activity to verify the intactness of lysosomes that were isolated from HEK293 cells based on anti-TMEM192 Lyso-IP. Our data showed an increase in the cathepsin D activity of lysosomal fraction when compared with whole cell fraction and Mock-IP fraction, an indication that the lysosomes are intact and viable.

Attachments



Cathepsin D assay fo...

28KB




Materials

Materials:

1. Cell lines

-  HEK293 **ATCC Catalog #CRL-1573** , RRID:CVCL_0045)


2. Media and Reagents

-  Cathepsin D Activity Assay Kit (Fluorometric) **Abcam Catalog #ab65302**

3. Equipment

- ClarioStar plate reader

4. Consumables




-  FLUTAC flat bottom black 96-well plate **greiner bio-one Catalog #655076**
- Standard 1ml and 200µl Pipette tips (Greiner bio-one. Catalog# 686271 and 685261 respectively).



Seeding cells and performing Lyso-IP with anti-TMEM192 beads



- 1 Seed HEK293 cells in 15cm plates and allow to reach 80-90% confluency.
- 2 Perform Lyso-IP (using anti-TMEM192 beads) and Mock-IP (using BSA coated beads) as previously described in [dx.doi.org/10.17504/protocols.io.x54v9yp51g3e/v1](https://doi.org/10.17504/protocols.io.x54v9yp51g3e/v1)

Preparing sample for Cathepsin D assay

- 3 Add  2 μg of protein from Lyso-IP and whole cell lysate into the wells FLUOTAC flat bottom black 96-well plate. This should be done in duplicate. 
- 4 Top up to  50 μL with lysis buffer provided in the kit.

Note

Due to little or no protein in the Mock-IP, use equal volume as Lyso-IP sample.

- 5 Prepare Blank sample in duplicate. This should contain only lysis buffer.
- 6 Prepare a reaction master mix for 9 wells:
 -  450 μL reaction buffer (from the kit) per well.
 -  18 μL substrate (from the kit)

Note

- Although there are 8 wells to be used, however make master mix for 9 wells to account for pipetting error.
- If performing assays for more samples/replicates, adjust master mix accordingly.









- 7 Add  52 μL of master mix into each well. Gently mix but avoid bubbles. Cover plate with foil to avoid light exposure.

Plate reading and analysis.

2h

- 8 Start the ClarioStar plate reading machine and initiate the software.
- 9 Set temperature to  37 $^{\circ}\text{C}$.
- 10 Set reading time to  00:05:00 for 24 cycles. This is total reading time of  02:00:00 .
- 11 Set reading wavelength to Ex/Em =  328 undetermined /  460 undetermined .
- 12 Highlight and name the virtual wells, ensuring they correspond with the orientation of the plate.
- 13 Set direction of plate reading.
- 14 Open the plate holder and insert plate. Remember to remove foil covering before inserting the plate into the equipment.
- 15 Close plate holder and run the program.
- 16 After the completion of the run, export data in excel format and analyse it using GraphPad Prism.




Figure 1: Cathepsin D activity showed that the purified lysosomes are intact and retain their content. After IP, Measure Cathepsin D activity from 2ug of protein obtained from lysosomal fraction and whole cell fraction while the Mock-IP serves as negative control. N=2