




Mar 13, 2021

# Chimeric Protein-LAG and Protein-LG sandwich ELISA

Angel A Justiz-Vaillant<sup>1</sup><sup>1</sup>University of the West Indies St. Augustine**1** Works for me [dx.doi.org/10.17504/protocols.io.btbdnii6](https://dx.doi.org/10.17504/protocols.io.btbdnii6)

Carbon

 Angel A Justiz-Vaillant  
University of the West Indies St. Augustine

SUBMIT TO PLOS ONE

## ABSTRACT

This ELISA was used to study the interactions between protein-LAG (PLAG) and protein-LG (SpLG) with different immunoglobulin preparations from mammalian species.

DOI

[dx.doi.org/10.17504/protocols.io.btbdnii6](https://dx.doi.org/10.17504/protocols.io.btbdnii6)

## PROTOCOL CITATION

Angel A Justiz-Vaillant 2021. Chimeric Protein-LAG and Protein-LG sandwich ELISA. **protocols.io**  
<https://dx.doi.org/10.17504/protocols.io.btbdnii6>

## LICENSE

————— This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

## CREATED

Mar 13, 2021

## LAST MODIFIED

Mar 13, 2021

## PROTOCOL INTEGER ID

48197

- 1 This ELISA was used to study the interactions between protein-LAG (PLAG) and protein-LG (SpLG) with different immunoglobulin preparations from mammalian species. The 96 well microtiter plate was coated overnight at 4°C with 2 µg/µl per well of PLAG in carbonate-bicarbonate buffer pH 9.6.
- 2 The plate was then treated with bovine serum albumin solution and washed 4X with PBS-Tween. 50 µl of immunoglobulins (1 mg/ml) is added and incubated for 1h at room temperature, and the microplate is rewashed 4X with PBS-Tween.
- 3 Then, 50 µL of peroxidase-labeled SpLG conjugate diluted 1:5000 in PBS-non-fat milk was added to each well and incubated for 1h at RT. The plate was washed 4X with PBS-Tween.

- 4 Then, 50  $\mu$ L of o-phenylenediamine solution (4 mg/mL) was added, and the plate was incubated for 15 min at RT in the dark. The reaction was stopped with 50  $\mu$ L of a 3M H<sub>2</sub>SO<sub>4</sub> solution.
- 5 The plate was visually assessed for color development and read on a microplate reader at 492 nm. A cut-off point was calculated as the mean of the optical density of the negative controls multiplied by two. The cut-off value was 0.34.