



Aug 15, 2020

# Binding properties of immunoglobulin-binding protein by double immunodiffusion (Ouchterlony) technique.

Angel A Justiz-Vaillant<sup>1</sup><sup>1</sup>University of the West Indies St. Augustine**1** Works for me [dx.doi.org/10.17504/protocols.io.bjsjknkn](https://dx.doi.org/10.17504/protocols.io.bjsjknkn)[University of the West Indies](#) [angel.vaillant@sta.uwi.edu](mailto:angel.vaillant@sta.uwi.edu)Angel Justiz-Vaillant  
University of the West Indies St. Augustine

DOI

[dx.doi.org/10.17504/protocols.io.bjsjknkn](https://dx.doi.org/10.17504/protocols.io.bjsjknkn)

## PROTOCOL CITATION

Angel A Justiz-Vaillant 2020. Binding properties of immunoglobulin-binding protein by double immunodiffusion (Ouchterlony) technique.. **protocols.io**  
<https://dx.doi.org/10.17504/protocols.io.bjsjknkn>

## LICENSE

This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

## CREATED

Aug 15, 2020

## LAST MODIFIED

Aug 15, 2020

## PROTOCOL INTEGER ID

40491

- 1 The binding of SpLAG, SpLG and SpAG with animal sera, avian IgY, and purified mammalian IgG are investigated by double immunodiffusion.
- 2 Briefly, 1% agarose gels are prepared and wells cut into the gel using a template.
- 3 Initially, aliquots of 25 µl each of the immunoglobulin-bacterial protein SpLAG, SpLG or SpAG at a concentration of one (1) µg/µl are applied to the centre well.
- 4 The peripheral wells are filled with 25 µl each of IgY (30 µg/µl) or mammalian IgG or sera.
- 5 The gels are incubated at RT for 48–72 hours.

- 6 After that the gels are examined for precipitin lines.
- 7 Human serum or human IgG are included as positive controls.
- 8 The positive results are taken as the presence of precipitin line/s and negative results, the absence of precipitin lines.
- 9