

Oct 02, 2024


READDI protocol: Crystallisation of CHIKV nsP3 macrodomain



Forked from [READDI protocol: Crystallisation of CHIKV nsP3 macrodomain](#)

DOI

dx.doi.org/10.17504/protocols.io.j8nlk8qzdl5r/v1



Jasmin Aschenbrenner^{1,2}, Peter Marples^{1,2}, Daren Fearon^{1,2}

¹Diamond Light Source; ²Research Complex at Harwell

ASAP Discovery



Peter Marples

Diamond Light Source

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.j8nlk8qzdl5r/v1

External link: <https://readdi-ac.org/>

Protocol Citation: Jasmin Aschenbrenner, Peter Marples, Daren Fearon 2024. READDI protocol: Crystallisation of CHIKV nsP3 macrodomain. [protocols.io https://dx.doi.org/10.17504/protocols.io.j8nlk8qzdl5r/v1](https://dx.doi.org/10.17504/protocols.io.j8nlk8qzdl5r/v1)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: October 02, 2024

Last Modified: October 02, 2024

Protocol Integer ID: 108825

Keywords: crystallisation, XChem, Diamond Light Source, i04-1, READDI, chikungunya virus, AViDD, Research complex at Harwell, CHIKV NS3

**Funders Acknowledgement:**

**National Institutes of
Health/National Institute Of
Allergy and Infectious
Diseases (NIH/NIAID)**
Grant ID: U19AI171292

Disclaimer

The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Acknowledgements:

Diamond Light Source Ltd, Harwell Science and Innovation Campus, Didcot OX11 0QX, UK
Research Complex at Harwell, Harwell Science and Innovation Campus, Didcot OX11 0FA, UK
Oxford Lab Technologies crystal shifter <https://doi.org/10.1107/S2059798320014114>

Abstract

The crystallization protocol and buffer conditions used to obtain reproducible Chikungunya Virus NS3 macrodomain crystals suitable for **XChem** fragment screening.

Guidelines

N/A

Materials


SwissCI 3 lens crystallization plates <https://swissci.com/product/3-lens-crystallisation-plate/> **Codes:**
Midi: UVXPO-3LENS 3W96T-PS 3W96T-UVP

Molecular Dimensions 'The BCS Screen Single Reagent' 2-44:

0.1 M Tris (pH 7.8), 0.1 M Potassium thiocyanate, 0.1 M Sodium bromide, 25 % v/v PEG Smear Broad, Catalog # MDSR-104-2-44

Purified CHIKV Mac protein (11 mg/mL) in 25 mM Tris-HCl (pH 7.5), 0.1 M NaCl, 5 % Glycerol.

Safety warnings

 Follow all handling warning for the chemicals used in the crystallisation screen composition.

Ethics statement

N/A

Equipment needed

- 1 **Formulatrix Rock Imager** (or incubator of choice)
SPT mosquito

Equipment

Mosquito HV

NAME

High Volume 16-Channel Robotic Liquid Handler

TYPE

SPT LabTech

BRAND

3097-01057

SKU

<https://www.sptlabtech.com/products/liquid-handling/mosquito-hv/>^{LINK}

P100 8 multi-channel pipette

SwissCI 3 lens plate

Crystallisation experiment

1d

- 2 **Prepare seed stock:**

17m 40s

Protocol



NAME

Diamond XChem Seeding Protocol

CREATED BY

Peter Marples

PREVIEW

1: 100 dilution  Sample seeds

- 3 **Protein and buffer requirements:**



21.6 µL



1M 11 mg/mL



Sample



🧪 2.88 mL Crystallisation screen

🧪 10.08 µL Sample Seeds, dilution 1:100

4 Crystallisation screen composition:

[M] 0.1 Molarity (M) Tris-NaOH 7.8

[M] 0.1 Molarity (M) Potassium thiocyanate

[M] 0.1 Molarity (M) Sodium bromide

25 % v/v PEG Smear Broad

Stock solutions used:

[M] 1 Molarity (M) Tris adjusted to 7.8 with NaOH

[M] 1 Molarity (M) Potassium thiocyanate

[M] 1 Molarity (M) Sodium bromide

50% v/v PEG Smear Broad

Note

The crystallisation screen can be stored in a duran bottle or aliquoted into 96 deep well block for easy dispensing into SwissCI 3 lens plates.

For long term storage keep the crystallisation screen in the fridge at 4°C.

5 Dispense 30 µL Crystallisation screen into SwissCI 3 lens plate reservoir wells using a 100 µl multi-channel pipette.

10m

Dispense 75 undetermined [M] 11 mg/mL Sample to each lens using the SPT mosquito.

Dispense 40 undetermined Crystallisation screen to each lens using the SPT mosquito.

Dispense 35 undetermined CHIKV Mac Seeds to each lens using the SPT mosquito.

Drop ratio: 15:8:7 ratio (75 nl Sample : 40 nl Crystallisation solution: 35 nl Seeds)

Final drop volume: 150 nl

6 Incubate at 20 °C for 24:00:00 h in Formulatrix Rock Imager.

1d

Imaging Schedule: The first images are taken after 12 h and the imaging schedule follows a Fibonacci sequence of days for further collections.

7

Expected result

The crystals reach their maximum size after 24-48 h.

Crystals typically form as single crystals at the bottom of the drop or on the drop-air interface.

Morphology: typically thin rectangles with pointed ends.


Size: ~100 μm in length and ~50 μm in width, depth of the crystals is ~10 μm

Appearance: glass shard.

Average resolution: 1.5 \AA

Space group: $P3_1$

Unit cell: 87 \AA , 87 \AA , 85 \AA
90.00°, 90.00°, 120.00°



An example of a drop containing CHIKV macrodomain crystals.

Data collection at Synchrotron



8 Diamond Light Source
Unattended Data Collection (UDC)
Data Collection Temperature: 100K
Detector: DECTRIS EIGER2 X 9M
Beamline: I04-1
Wavelength: 0.9212 Å
Resolution (Å): 1.64
Beam Size (µm): 60 X 50
Number of images: 3600
Oscillation: 0.10°
Exposure (s): 0.0020
Transmission (%): 100
Flux (ph/s): 3.80e+12

Protocol references

Identifying novel chemical matter against the Chikungunya virus nsP3 macrodomain through crystallographic fragment screening, <https://doi.org/10.1101/2024.08.23.609196>