

MAR 17, 2023

Sequencing of construct

In 6 collections

Pascale Baden¹, michela.deleidi¹

¹German Center for Neurodegenerative Diseases (DZNE), Tübingen, 72076 Germany



Hariam Raji

ABSTRACT

This protocol describes sequencing of construct.

ATTACHMENTS

404-874.docx

MATERIALS

Materials

Fw primer (CMV FW)
Rv primer (BGH RV)
sodium acetate (Carl Roth)
EtOH (VWR Chemicals BDH Prolabo)
Terminator Sequencing buffer
MilliQ-H₂O

Hi-Di Formamide (Applied Biosystems)

⊠ GoTaq(R) DNA Polymerase, 500u **Promega Catalog #M3005**

Ø dNTP Set 100 mM Solutions Thermo Fisher Scientific Catalog #R0182

BigDye[™] Terminator v3.1 Cycle Sequencing Kit **Thermo Fisher Catalog**#4337455

OPEN ACCESS

יוסם

dx.doi.org/10.17504/protocol s.io.bp2l617ydvqe/v1

Protocol Citation: Pascale Baden, michela.deleidi 2023. Sequencing of construct. **protocols.io**

https://dx.doi.org/10.17504/protocols.io.bp2l617ydvqe/v1

License: This is an open access protocol distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working We use this protocol and it's working

Created: Apr 26, 2022

Last Modified: Mar 17, 2023

PROTOCOL integer ID: 61424

Keywords: Sequencing-PCR, Sodium acetate precipitation, Load Sequencing-plate

PCR to amplify region of interest



A	В
Master mix	1x
H2O	13.3
5x Colorless Reaction Buffer (Promega, #M3005)	4
10mM dNTPs (ThermoFisher, #R0182)	0.4
Fw primer (CMV FW)	0.6
Rv primer (BGH RV)	0.6
GoTaq Polymerase (Promega, #M3005)	0.1
Σ	19 µl

Mix \perp 19 μ L MM with \perp 1 μ L DNA (\perp 50 μ C).

PCR Program

2

94°C -> 5 min 94°C -> 30 sec 60°C* -> 30 sec 72°C -> 1 min 94°C -> 30 sec 50°C -> 30 sec 72°C -> 1 min 72°C -> 7 min 4°C -> endless * -1°C /cycle

Sodium acetate precipitation

3

Add \pm 50 μ L sodium acetate (\pm 1 mL [M] 3 Molarity (M) Na Acetate (Carl Roth) + \pm 24 mL 100% EtOH (VWR Chemicals BDH Prolabo)) to \pm 20 μ L PCR product.

15m



- **5** Remove supernatant (pat plate on paper).
- 6 Add Δ 100 μL 70% EtOH onto pellet.



7 Centrifuge at 3200 rcf, 4°C, 00:15:00



- **8** Remove supernatant (pat plate on paper).
- 9 Add \angle 100 μ L 70% EtOH onto pellet.



10 Centrifuge at 3200 rcf, 4°C, 00:15:00



11 Remove supernatant (pat plate on paper).

15m



remove EtOH.

13

Add \pm 15 μL MilliQ-H2O to pellet and vortex 15-20 min (speed 0-1).



Sequencing-PCR

14



A	В
	1x sample
H20	5.3 µl
5x Terminator Sequencing buffer	3.3 µl
BigDye v3.1 (ThermoFisher, #4337455)	1.4 µl
Σ	10 μΙ

Add 🗸 5 µL MM.

15

Add \pm 4 μ L DNA from step 13.



Add 🗸 1 µL primer FW or RV.



Sequencing program

Sodium acetate precipitation

18 Add \angle 25 μ L sodium acetate to \angle 10 μ L PCR product from.



19 Mix well and centrifuge at 3200 rcf, 4°C, 00:45:00

45m



- 20 Remove supernatant (pat plate on paper).
- 21 Add \pm 100 μ L 70% EtOH onto pellet.



22 Centrifuge at 3200 rcf, 4°C, 00:15:00

15m

- *****
- Remove supernatant (pat plate on paper).

Add Δ 100 μL 70% EtOH onto pellet.



25 Centrifuge at 3200 rcf, 4°C, 00:15:00

15m

- *****
- Remove supernatant (pat plate on paper).
- Centrifuge upside down max. 600 rcf, 4°C, 00:01:00 (top of sample down on tissue paper) to remove EtOH.

1m

- **®**
 - Add Δ 15 µL MilliQ-H₂O to pellet and vortex 15-20min (speed 0-1).



Note

Note: COVER! Light sensitive!

Load Sequencing-plate

29 Add 🗸 10 µL Hi-Di Formamide (Applied Biosystems).



30 A 7 µL DNA after purification (Step 28).

31 Store in 4 °C until sequencing.