



FEB 14, 2023

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.eq2ly776plx9/v1

Protocol Citation: Karine Lapouge, Rozemarijn Kleinendorst, Kasit Chatsirisupachai, Nikolay Dobrev, Kim Remans, Arnaud Krebs 2023. Expression, purification and characterization of the GpC methyltransferase M.CviPI.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.eq2ly776plx9/v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Nov 02, 2022

Last Modified: Feb 14, 2023


PROTOCOL integer ID:
 72195

🌐 Expression, purification and characterization of the GpC methyltransferase M.CviPI

Karine Lapouge¹, Rozemarijn Kleinendorst²,
 Kasit Chatsirisupachai², Nikolay Dobrev¹, Kim Remans¹,
 Arnaud Krebs²

¹European Molecular Biology Laboratory (EMBL), Protein Expression and Purification Core Facility, 69117, Heidelberg, Germany;

²European Molecular Biology Laboratory (EMBL), Genome Biology unit, Meyerhofstraße 1, 69117 Heidelberg. 2European Molecular Biology Laboratory (EMBL)



Arnaud Krebs

ABSTRACT

Methylation footprinting can be used to map protein-DNA contacts at the resolution of individual DNA molecules¹. Enzymes with various nucleotide specificities have been successfully used to footprint genomes including GpC, CpG and A methyltransferases²⁻⁷. Among these M.CviPI methylate DNA in GpC context, that is distinct from CpGs that are endogenously methylated in mammals. This feature has been leveraged to profile nucleosome occupancy^{8,9}; the binding of General Transcription Factors and RNA Pol II⁶; the co-occupancy of Transcription Factors (TFs)¹⁰ and the relation between TF binding and endogenous DNA methylation¹¹. Here, we present a protocol for the production and purification of M.CviPI in *E. coli*. Our protocol routinely yields milligrams of protein at a quality and a concentration compatible with DNA footprinting applications. We characterize the purity and the activity of the purified enzyme, providing a benchmark for future production.

ATTACHMENTS

[pBAD_HisMBP3C-McviPI.dna](#)

[Protocol_McviPI_purification.pdf](#)