



APR 04, 2024

🌐 Generation of induced neurons from human induced pluripotent stem cells.

Prarthana Gowda¹, Zhiping Pang², Mahmoud ElAchwah², Sol Diaz de Leon Guerrero²

¹Rutgers; ²Rutgers University



Prarthana Gowda
Rutgers

OPEN  ACCESS



DOI:

dx.doi.org/10.17504/protocols.io.e6nvwdbxzlmk/v1

Protocol Citation: Prarthana Gowda, Zhiping Pang, Mahmoud ElAchwah, Sol Diaz de Leon Guerrero 2024. Generation of induced neurons from human induced pluripotent stem cells. . **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.e6nvwdbxzlmk/v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Sep 20, 2023


ABSTRACT

Last Modified: Apr 04, 2024

PROTOCOL integer ID: 88114

Funders Acknowledgement:

NIH NIMH Assay and Data Generation Center (ADGC) for the Model of iPSC-derived Neurons for NPD (MiNND)
Grant ID: RM1MH133065-01



Overview of induced neuron protocol for the generation of excitatory and inhibitory neurons.

Protocol for the generation of Ngn2 (Excitatory) and Ascl1/Dlx2 (Inhibitory) induced neurons from human induced pluripotent stem cells (iPSCs) and co-culture of excitatory and inhibitory neurons on mouse glia (Protocols modified from Wang et al., 2022; Yang et al., 2017; Zhang et al., 2013).

iPSCs are maintained in 35mm plates or 6 well plates in mTESR+ media (<https://dx.doi.org/10.17504/protocols.io.ewov1qd5ogr2/v1>).

Lentiviral vectors were generated by transfecting HEK293T cells with lentivirus packaging plasmids (pMDLg/pRRE, VsVG and pRSV-REV) with the desired vectors as previously described (Pang et al., 2011) using lipofectamine 3000. The following plasmids were used: pMDLg/pRRE (Addgene 12251), pRSV-Rev (Addgene #12253), pCMV-VSV-G (Addgene #8454), FUW-M2rtTA (Addgene #20342), FUW-TetO-Ngn2-P2A-puromycin (Addgene #52047), FUW-TetO-Ascl1-T2A-puromycin (Addgene #97329), FUW-TetO-Dlx2-IRES-hygromycin (Addgene #97330). Lentiviral particles were collected in mTESR+ media and stored at -80°C until further use.

Induced neurons are generated by transducing iPSCs with the necessary genes using the lentiviral vectors to induce the expression of different transcription factors with doxycycline: rtTA + Ngn2 for excitatory neurons, rtTA + Ascl1 + Dlx2 for inhibitory neurons. Induced neurons are plated into 96 well plates (18×10^3 cells) or in 12 well plates (1×10^6 cells) with mouse glia after 5 days of induction. Primary mouse glia was obtained from postnatal day 0-2 mice cortex kept in DMEM 10% FBS 1% pen/strep, glia is used for plating after the second or third passage (P2-P3). CEPT cocktail was used during iPSC and neuronal plating to enhance cell viability (Chen et al., 2021). Induced neurons are cultured for 30-35 days before analysis.

PROTOCOL REFERENCES

Chen, Y., Tristan, C.A., Chen, L., Jovanovic, V.M., Malley, C., Chu, P.H., Ryu, S., Deng, T., Ormanoglu, P., Tao, D., et al. (2021). A versatile polypharmacology platform promotes cytoprotection and viability of human pluripotent and differentiated cells (Springer US).

Pang, Z.P., Yang, N., Vierbuchen, T., Ostermeier, A., Fuentes, D.R., Yang, T.Q., Citri, A., Sebastiano, V., Marro, S., Südhof, T.C., et al. (2011). Induction of human neuronal cells by defined transcription factors. *Nature* 476, 220–223. <https://doi.org/10.1038/nature10202>.

Wang, L., Mirabella, V.R., Dai, R., Su, X., Xu, R., Jadali, A., Bernabucci, M., Singh, I., Chen, Y., Tian, J., et al. (2022). Analyses of the autism-associated neuroligin-3 R451C mutation in human neurons reveal a gain-of-function synaptic mechanism. *Mol. Psychiatry* 1–16. <https://doi.org/10.1038/s41380-022-01834-x>.

Yang, N., Chanda, S., Marro, S., Ng, Y.H., Janas, J.A., Haag, D., Ang, C.E., Tang, Y., Flores, Q., Mall, M., et al. (2017). Generation of pure GABAergic neurons by transcription factor programming. *Nat. Methods* 14, 621–628. <https://doi.org/10.1038/nmeth.4291>

Zhang, Y., Pak, C.H., Han, Y., Ahlenius, H., Zhang, Z., Chanda, S., Marro, S., Patzke, C., Acuna, C., Covy, J., et al. (2013). Rapid single-step induction of functional neurons from human pluripotent stem cells. *Neuron* 78, 785–798. <https://doi.org/10.1016/j.neuron.2013.05.029>.

MATERIALS

Reagent	Vendor	Catalog Number	Stock concentration	Working concentration
Accutase	innovative cell technologies inc	AT-104	1x	1x
AraC	C1768	C1768	8mM	2-4 μ M
B27	Gibco	17504044	50x	1x
BDNF	pepro tech	10781-164	250 μ g	10ng/mL
CET/P	medchem express, selleck chem, R&D systems sigma-aldrich	HY-15392, S7775, 5284, P8483	C (50 μ M), E (5000 μ M), P(1x), T (0.7mM)	C (50nM), E (5uM), P(1x), T (0.7 μ M)
DMEM	Gibco	11995-065	1x	1X
Doxycycline	MP biomedical	198955	2mg/mL	2 μ g/mL
dPBS	SAFC	D8537	1x	1x
FBS	r&d systems	S11550	100%	1x
GDNF	Pepro tech	10781-226	200 μ g	10ng/mL
glutamax	gibco	35050061	100x	1x
hygromycin	sigma-aldrich	H9773	50mg/mL	100 μ g/mL
matrigel	Cornig	354234		
MEM	gibco	51200-038	1x	1x
mTESR+	Stem cell technology	100-0275	1x	1x + supplement
neurobasal	gibco	21103-049	1x	1x
NT3	pepro tech	10781-174	250 μ g	10ng/mL
penicillin/streptomycin	thermo-fisher	15070-63	100x	1x
puromycin	sigma-aldrich	P8833	1mg/mL	1-2 μ g/mL
Primocin	InvivoGen	ant-pm-1	50mg/mL	100 μ g/mL
0.05% trypsin EDTA	gibco	25300-054	1x	1x
12 well plate	falcon	353043	1mL/well	4cm ²

	Reagent	Vendor	Catalog Number	Stock concentration	Working concentration
	6 well plate	biofil	2304117-074-F	2mL/well	9cm ²
	96well plate	greiner	655090	200uL/well	0.32cm ²
	24well plate	biofil	230606-076-F	1mL/well	2cm ²

Media preparation

Neurobasal (neuronal for induction, selection , recovery)- 500mL neurobasal, 10mL B27 (0.5mM), 5mL glutamax supplement

Neurobasal + FBS (neuronal media for maintenance) - 500mL neurobasal, 10mL B27 (0.5mM), 5mL glutamax supplement, 25mL FBS (5%)

mTeSR+ (iPSC maintenance and infection) - 400uL mTeSR, 100mL mTeSR supplement, 1mL Primocin

DMEM - 500mL DMEM, 50mL FBS (10%), 5mL penecillin/streptomycin (1%)

**all media filtered during preparation.*















Infection of iPSCs - day 0





3h

- 1 Follow the below protocols for excitatory and inhibitory neuron generation.




STEP CASE

Excitatory neurons 46 steps



- 2 Warm up  1.5 mL of mTESR+ per cell line
 Warm up  2.5 mL MEM per cell line
 Warm up  1 mL of accutase per cell line
 Thaw required volumes of NGN2 and RTTA virus on ice
 Coat and incubate 12 well plates (1 well per line) with  500 μ L of Matrigel per well for at least
 01:00:00
- 3 Aspirate old media from confluent iPSC plate. Always use different tips to avoid contamination between the lines.
- 4 Wash with  500 μ L MEM
- 5 Add  500 μ L accutase, incubate at  37 $^{\circ}$ C and 5% CO₂ for  00:06:00 6m
- 6 Make sure all the cell colonies are suspended, transfer all cell from each well to  2 mL of MEM and 5m
 centrifuge for  00:05:00 at 1000rpm at  23 $^{\circ}$ C .
- 7 Add 1:1000 of CET/P ( 1 μ L of CET and  1 μ L of P per mL of media) to mTESR+, mix well

- 8 Add NGN2 and RTTA virus (1:1) to the  500 μ L mTESR+ with CET/P per line, mix well (discard all tips and tubes in 10% bleach)
- 9 Aspirate excess Matrigel from well
- 10 Add  500 μ L of virus+mTESR+ with CET/P to a labelled well.
- 11 Aspirate out the supernatant, and resuspend each line of cells will  1 mL mTESR+, make sure cells are resuspended well to ensure cells do not form large colonies for better infection.
- 12 Use 1:1 dilution of trypan blue to cell to count cells using an automatic cell counter. For a 12 well plate seed $\sim 1 \times 10^5$ - 1.25×10^5 cells. For a 6 well plate seed $\sim 2 \times 10^5$ - 2.5×10^5 cells. Add an appropriate volume of cell suspension to the labelled well(s), incubate at  37 °C and 5% CO₂


Induction - day 1




- 13 Warm up  1 mL of Neurobasal+B27+Glutamax per well of a 12 well plate.
- 14 Prepare induction media
 Add 1:1000 of Doxycycline ( 1 μ L). *2 μ g/mL stock solution, 2 μ g/mL working solution*
 Add 1:1000 CET/P ( 1 μ L of each). *2 μ g/mL stock solution, 2 μ g/mL working solution*

15 Remove media from the 12 well plate into 10% bleach (including all tips and tubes)



16 Add  1 mL of induction media (NB+Glut+B27+dox+CET/P) to each well. Incubate at  37 °C and 5% CO₂

Selection - day2&3

17 Prepare selection media (NB+Glut+B27+puro+dox). Warm up  1 mL neurobasal+glutamax+B27 per well of a 12 well plate



18 Add 1:1000 of Doxycycline( 1 µL)
Add 1:1000 - 1:500 of puromycin ( 1 µL -  2 µL) - depending on the number of iPSCs

19 Aspirate out old media from the 12 well plate












20 Add  1 mL of selection media (NB+Glut+B27+puro+dox) to each well, Incubate at  37 °C and 5% CO₂








Recovery- day 4

21 Take out  1 mL of Neurobasal+glutamax+B27 per well of a 12 well plate

- 22 Add 1:1000 of Doxycycline ( 12 μ L)
- 23 Aspirate out old media from 12 well plate
- 24 Add  1 mL of recovery media (NB+Glut+b27+dox) to each well, Incubate at 37°C and 5% CO₂







Glia plating - day 4


- 25 Coat 96 well plates for sensor (5 wells per line) or high content imaging (8 wells per line) experiments with  100 μ L matrigel, 12 well plate for RNAseq experiment (1 well per line) with  500 μ L materiel for  01:00:00 1h
- 26 Warm up  5 mL trypsin,  10 mL of NB+B27+Glutmax+5% FBS (plating media),  3 mL of NB+B27+Glutmax+5% FBS (resuspension), and  5 mL DMEM+10%FBS+1%penecillin-streptomycin
500mL neurobasal, 10mL B27 (0.5mM), 5mL glutamax supplement, 25mL FBS
- 27 Select a confluent plate of P2/P3 mouse glia
- 28 Aspirate out old media from the glia, wash with  5 mL dPBS
- 29 Add  5 mL of trypsin, incubate for  00:05:00 at  37 °C and 5% CO₂ 5m



- 30** Make sure all the cells are lifted and transfer the trypsin suspended cell to  5 mL of DMEM+10%FBS+1%p/s, centrifuge for  00:05:00 at 1000rpm at  23 °C . 5m
- 31** Aspirate out supernatant, and resuspend cells with  3 mL NB+B27+Glutmax+5% FBS
- 32** Count cell in  3 mL suspension, calculate volume for 8×10^3 - 1.2×10^3 cells per well of a 96well plate or 2.5×10^3 - 4×10^3 cells per well of a 12well plate.
(number of glia seeded depends on if the cultures are P2 or P3 and how old the cultures are at the time of plating)
- 33** Aspirate excess Matrigel from the wells.
- 34** Add the calculated volume of cell suspension required for plating in 100uL per well of 96well plate and  750 μ L per well of a 12 well plate. Add cell volume to plating media. Mix well, add to well. Incubate at  37 °C and 5% CO₂




iN plating - day5


5m



- 35** Warm up  10 mL of Neurobasal+Glutamax+B27+5%FBS (plating media) per 96well plate ( 100 μ L per well) or 12 well plate  750 μ L per well).
Warm up  500 μ L of Neurobasal+Glutamax+B27+5%FBS (plating media) per cell line for resuspension.
Warm up  500 μ L of accutase per well
Warm up  3 mL of MEM for each well
- 36** Aspirate out old media from excitatory iNs well



- 37 Wash with  500 μ L of MEM

- 38 Add  500 μ L of accutase to each well, Incubate at 37°C and 5% CO₂ for  00:06:00 6m

- 39 Transfer suspended cell into  2 mL of MEM in 15mL falcon tube, centrifuge for  00:05:00 at 5m
1000rpm at  23 °C




- 40 Resuspend each line in  500 μ L

- 41 To count cells, mix  10 μ L trypan blue with  10 μ L cell suspension, add 10 μ l of diluted cells to one sides of the cell counter.




- 42 Calculate volume for 12x10³ cells per well of a 96 well plate or 8.4x10⁵ cells per well of a 12 well plate, add calculated volume to  500 μ L plating media for each well. Make sure to label each well with the condition. Mix well. Incubate at  37 °C and 5% CO₂

- 43 Excitatory and inhibitory cells are co-cultured for all experiments mentioned above, check inhibitory neuron protocol for more details.





Day6

- 44** Change media to neuronal media -  100 μ L for a 96 well plate or  750 μ L of NB+Glut+B27+5%FBS + Factors per well of a 12 well plate
- Factors: 1:1000 Doxycycline
- | | |
|---------------------------|---|
| 1:1000 100ug/mL GDNF | 10ng/mL working solution (1:10 dilution in dPBS) |
| 1:1000 100ug/mLBDNF. | 10ng/mL working solution |
| 1:1000 100ug/mL NT3. | 10ng/mL working solution |
| 1:2000 or 1:4000 8mM AraC | 2uM-4uM working solution (based on glia density, only first feeding- either day6 or day8) |
| 1:500uL of Primocin. | 100ug/mL working solution (only first feeding - day6) |
- Incubate at  37 °C and 5% CO₂

Day8

- 45** Add  100 μ L or  750 μ L of neuronal media with factors (except for doxycycline and primocin) per well of a 96 well plate and a 12well plate, respectively.
- NB+Glut+B27+5%FBS + Factors
- Factors: 1:1000 GDNF
- 1:1000 BDNF
- 1:1000 NT3
- 1:2000 or 1:4000 AraC (based on glia density, only first feeding- either day6 or day8)
- Incubate at  37 °C and 5% CO₂

Day 13, Day18, Day23, Day28, Day 33....

- 46** Discard half of the old media, replace with  100 μ L or  750 μ L media every 5 days (make sure outside wells are not evaporating media faster) per well of a 96well plate and 12 well plate, respectively.
-  1 mL NB+Glut+B27+5%FBS + Factors
- Factors: 1:1000 GDFN
- 1:1000 BDFN
- 1:1000 NT3
- Incubate at  37 °C and 5% CO₂
- 47** Maintain cultures for 30-35days

