



JAN 31, 2024

Direct acid extraction

Sigrid Verhelst¹, Laura Corveleyn¹

¹ProGenTomics, Laboratory of Pharmaceutical Biotechnology, Ghent University, Ghent, Belgium



Laura Corveleyn

ProGenTomics, Laboratory of Pharmaceutical Biotechnology, Gh...

DISCLAIMER

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to [protocols.io](https://dx.doi.org/10.17504/protocols.io.bp2l6x6odlqe/v1) is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with [protocols.io](https://dx.doi.org/10.17504/protocols.io.bp2l6x6odlqe/v1), can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

ABSTRACT

Protocol to extract histones from cells using direct acid extraction with the purpose of label-free LC-MS/MS measurement.

OPEN ACCESS



DOI:

dx.doi.org/10.17504/protocols.io.bp2l6x6odlqe/v1

Protocol Citation: Sigrid Verhelst, Laura Corveleyn 2024. Direct acid extraction.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.bp2l6x6odlqe/v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: Jan 31, 2024

Last Modified: Jan 31, 2024

PROTOCOL integer ID: 94448

Funders Acknowledgement:

BOF

Grant ID: Mandate 01D23313

MATERIALS

Protein lobind Eppendorf tubes

Thermoshaker

Cooled centrifuge (4°C)

Ice

MilliQ water

12 M HCL (dilute to 0.4N HCl: for 300 mL: 10 mL 12M (37%) stock + 290 mL milliQ water)

TCA powder (make 100% TCA solution)

Ice-cold acetone

- 1 Resuspend the pellet in 0,4 N HCl by soft pipetting until no clumps are left in solution (**125 µL for 1x10⁶ cells**)

Note

If necessary, vortex softly

- 2 Incubate **4h** in acid on rotator **4°C** to promote lysis of nuclei and solubilization of histones.

- 3 Spin for 10 minutes at **4°C, 16000 g**.

- 4 Transfer supernatant to new Eppendorfs.

Note

Histones are present in the acid since they are alkaline proteins

- 5 Add, drop by drop (very slowly and in the middle of the eppendorf), TCA (**final concentration 33%: 61.25µl for 1x10⁶**) to promote precipitation of histones and invert the tube several times.

6 Incubate on ice for 30 minutes.

7 Spin for 10 min at 4°C, **16000g** to pellet the histones.

Note

Important: Place all the eppendorfs in the same position => makes it easier to know where the pellet is localized

8 Remove the supernatant.

Note

Be careful, the pellet is not always visible, smear on wall.

9 Add ± 150 μ L ice-cold acetone (don't resuspend the pellet) to remove TCA.

10 Spin for 5 min at 4°C, **16000g**.

11 Remove the supernatant.

12 Add ± 150 μ L cold acetone (don't resuspend the pellet) to remove TCA. If not all the acid is removed it will destroy the SDS-gel.

- 13 Spin for 5 min at 4°C, **16000g**.
- 14 Remove the supernatant.
- 15 Dry at room temperature for 30 minutes (until no acetone left).
- 16 Resuspend in milliQ water (**50 µL for 1x10⁶ cells**).
- 17 Transfer 20 µL (from 400.000 cells) to a new Eppendorf tube for gel.
- 18 Vacuum dry samples.