

Jun 21, 2024

DESS (DMSO/EDTA/NACL) Protocol

DOI

dx.doi.org/10.17504/protocols.io.5jyl822n9l2w/v1

Mirayana Marcelino Barros¹, Tiago Pereira², Alejandro De Santiago Perez¹, Hunter Powell³, Jenna Brown⁴

¹University of Georgia; ²Department of Marine Sciences, University of Georgia; ³UGA; ⁴University of Georgia- Holly Bik



Jenna Brown

University of Georgia- Holly Bik

OPEN ACCESS



DOI: dx.doi.org/10.17504/protocols.io.5jyl822n9l2w/v1

Protocol Citation: Mirayana Marcelino Barros, Tiago Pereira, Alejandro De Santiago Perez, Hunter Powell, Jenna Brown 2024. DESS (DMSO/EDTA/NACL) Protocol. protocols.io https://dx.doi.org/10.17504/protocols.io.5jyl822n9l2w/v1

License: This is an open access protocol distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working We use this protocol and it's

working

Created: May 29, 2024

Last Modified: June 21, 2024

Protocol Integer ID: 100828

Funders Acknowledgement:

Holly Bik

Abstract

DESS (DMSO/EDTA/NACL) protocol instructing how to make the solution for the preservation of samples.



Make 2L stock solution of 0.5M EDTA; if stock is already made, proceed to step 4.

- 1 Combine 372.24 g **disodium EDTA** and 500 mL deionized water.

 Tip: Disodium EDTA is labeled PINK in the Bik Lab.
- Add enough **5M NaOH** to the solution to bring it to a pH of \bigcirc 8.0 Tip: NaOH is labeled GREEN in the Bik Lab.

 Tip: In some cases, this can be as much as 500 mL. The EDTA will begin to dissolve around a pH of \bigcirc 7.0 .
- Bring final volume to 🚨 2 L with deionized water once all disodium EDTA has dissolved.

Prepare 2L DESS from 0.5M EDTA.

- Combine 1 L 0.5M disodium EDTA, 400 mL Dimethyl Sulfoxide/DMSO, and deionized water in a 4000 mL Erlenmeyer flask.

 Tip: 0.5M disodium EDTA is labeled YELLOW and Dimethyl Sulfoxide/DMSO is labeled ORANGE in the Bik Lab.
- 5 Put magnetic stir bars in the solution and place flask onto a hot plate. Turn the magnetic stirring function on and leave it at room temperature.
- Add enough **NaCl** to saturate the solution: Add NaCl in 100g increments until 300g is reached, and then proceed to adding smaller amounts until salt is no longer dissolving. Solution should saturate with the addition of ~300-400g NaCl.

Tip: NaCl is labeled RED in the Bik Lab.

Tip: This step can take hours: BE PATIENT. There may be leftover salt residue once the solution is fully saturated.