



Aug 23, 2020

ELISA for quantification of human properdin in serum or plasma.

Angel A Justiz-Vaillant¹, Belkis Ferrer-Cosme²¹University of the West Indies St. Augustine; ²Saturnino Lora Torres' Provincial Teaching Clinical Surgical Hospital. Cuba*In Development*dx.doi.org/10.17504/protocols.io.bj7zkrp6

University of the West Indies

angel.vaillant@sta.uwi.eduAngel Justiz-Vaillant
University of the West Indies St. Augustine

DOI

dx.doi.org/10.17504/protocols.io.bj7zkrp6

PROTOCOL CITATION

Angel A Justiz-Vaillant, Belkis Ferrer-Cosme 2020. ELISA for quantification of human properdin in serum or plasma.. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bj7zkrp6>

LICENSE

————— This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Aug 23, 2020

LAST MODIFIED

Aug 23, 2020

PROTOCOL INTEGER ID

40921

- 1 An anti-human properdin coating antibody is adsorbed onto the microwells by incubation overnight at 4°C with carbonate-bicarbonate buffer.
- 2 Add 50 µl of human serum or plasma. Human properdin present in the serum or plasma binds to antibodies adsorbed into the microwells.
- 3 The microplate is blocked with 3% non-fat milk-PBS buffer and later wash to remove unbound proteins.
- 4 Fifty (50) µl of biotin-conjugated anti-properdin antibody is added. The optimal dilution must be investigated.
- 5 The microplate is rewashed with PBS-Tween 20 buffer, pH 7.4.

- 6 One hundred μl of streptavidin-HRP conjugate is added and it binds to the biotin-conjugated anti-properdin antibody. The optimal dilution of this conjugate must be investigated.
- 7 The plate is washed following incubation to remove the unbound Streptavidin-HRP.
- 8 Add 100 μl of 3,3',5,5'- tetramethylbenzidine (TMB; Sigma-Aldrich) into each well.
- 9 Incubate the microwells in the dark for 20 min.
- 10 A colored product is formed in proportion to the quantity of human properdin present in the sample or standard.
- 11 The reaction is terminated by addition of 100 μl 3M H_2SO_4 and the absorbance is measured at 450 nm.
- 12 A standard curve is made from 7 human properdin standard dilutions and the human properdin sample concentration is determined.
- 13 For better results place the microplate on a microplate shaker in every incubation.