



Feb 07, 2020

Table 2. Multilocus genotyping results of the 24 *Giardia*-positive samples successfully genotyped at least at one of the three loci investigated. Shushtar County (Iran), 2017–2018. [↗](#)

PLOS One

Abdollah Rafiei¹, Raheleh Baghlaninezhad¹, Pamela C. Köster², Begoña Bailo², Marta Hernández de Mingo², David Carmena², Esmat Panabad³, Molouk Beiromvand¹

¹Infectious and Tropical Diseases Research Center, Health Research Institute, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran, ²Parasitology Reference and Research Laboratory, National Centre for Microbiology, Majadahonda, Madrid, Spain, ³Department of Parasitology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

1 Works for me dx.doi.org/10.17504/protocols.io.bapgidjw

Molouk Beiromvand

EXTERNAL LINK

<https://doi.org/10.1371/journal.pone.0228317>

THIS DOCUMENT ACCOMPANIES THE FOLLOWING PUBLICATION

Rafiei A, Baghlaninezhad R, Köster PC, Bailo B, Mingo MHD, Carmena D, Panabad E, Beiromvand M (2020) Multilocus genotyping of *Giardia duodenalis* in Southwestern Iran. A community survey. PLoS ONE 15(2): e0228317. doi: [10.1371/journal.pone.0228317](https://doi.org/10.1371/journal.pone.0228317)



This is an open access document distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited