




DEC 04, 2023

# eDNA extraction from buffer EtOH 100% using DNeasy Blood and Tissue Kit (QIAGEN)

delphine.vanhaecke<sup>1</sup><sup>1</sup>Universidad de Aysén

MOBI



delphine.vanhaecke

OPEN  ACCESS

## ABSTRACT

This is a protocol for extracting eDNA from the buffer EtOH 100% in which Millipore filters are stored.

### DOI:

[dx.doi.org/10.17504/protocols.io.3byl4qrojvo5/v1](https://dx.doi.org/10.17504/protocols.io.3byl4qrojvo5/v1)

**Protocol Citation:** delphine.vanhaecke 2023. eDNA extraction from buffer EtOH 100% using DNeasy Blood and Tissue Kit (QIAGEN).

**protocols.io**

<https://dx.doi.org/10.17504/protocols.io.3byl4qrojvo5/v1>

**License:** This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working  
We use this protocol and it's working

**Created:** Nov 14, 2023

**Last Modified:** Dec 04,  
2023

**PROTOCOL integer ID:**  
90941

**Funders**

**Acknowledgement:**

Gobierno Regional de Aysén  
Grant ID: FIC 2022 código BIP  
40049315-0

- 1 Clean the laminar flow hood and micropipettes with DNAZAP (solution 1 and 2) and rinse with ddH<sub>2</sub>O.
- 2 Place disposable gloves, sleeves and 1.5ml sterilized tubes (as many as the number of samples to process + 1 negative extraction control) in the laminar flow hood and submit to UV light for 10 minutes.
- 3 In the meantime, sterilize the tubes containing the filters with bleach (10%). Let dry and then clean with Ethanol 70%.
- 4 Beadbeat the tube with the filter (stored in EtOH 99.9%) for 20s at 4.0M/S. Let tubes rest for 5min then beadbeat again.
- 5 In the flowhood, remove the EtOH buffer from the tube and pipet it in a new 1.5mL sterilized LoBind tube
- 6 Centrifuge the LoBind tube with EtOH for 60 minutes at 8000rpm and 6°C (check if pellet has formed at bottom).
- 7 Refill the tubes with the filter with EtOH and store at -80°C.

- 8** In the flowhood, remove the supernatant and let the pellet dry with the fan on for 30 min.
- 9** Add 20µl proteinase K, 200µl PBS 1x, 200µl buffer AL to each pellet.  
Freshly prepare 5ml of PBS 1x from PBS10X (mix 500µl PBS10x with 4500µl nuclease free water).
- 10** Vortex well.
- 11** Incubate samples at 56°C for two hours in a thermoshaker at 600rpm.
- 12** Vortex 15s.
- 13** In the flowhood, add 200 µl EtOH 100%.
- 14** Mix by vortexing.
- 15** Pipet the mixture (650 µl) into the DNeasy Mini spin column placed in a 2 ml collection tube. Centrifuge at 8000 rpm for 1 min. Discard flow- through and collection tube.

- 16** Place the DNeasy Mini spin column in a new 2 ml collection tube, add 600 µl Buffer AW1, and centrifuge for 1 min at 8000 rpm. Discard flow-through and collection tube.
- 17** Place the DNeasy Mini spin column in a new 2 ml collection tube, add 600 µl Buffer AW2, and centrifuge for 3 min at 14,000 rpm to dry the Dneasy membrane. Discard flow-through and collection tube.
- 18** Place the DNeasy Mini spin column in a clean 1.5 ml microcentrifuge LoBIND tube, and pipet 75 µl Buffer AE directly onto the DNeasy membrane. Incubate at room temperature for 1 min, and then centrifuge for 1 min at 8000 rpm to elute.
- 19** Repeat step 18 in a new 1.5 ml microcentrifuge LoBIND tube for a 2nd elution.
- 20** Discard spin column and store eDNA samples at -20C.