

6



Apr 18, 2022

Second Balance beam assay to detect motor phenotypes in mice

William Hancock-Cerutti^{1,2,3,4,5}, Marianna Leonzino^{1,2,3,5}, Pietro De Camilli^{1,2,3,5}

¹Departments of Neuroscience and of Cell Biology, Yale University School of Medicine, New Haven, Connecticut 06510, USA;

²Howard Hughes Medical Institute;

³Program in Cellular Neuroscience, Neurodegeneration and Repair, Yale University School of Medici ne, New Haven, Connecticut 06510, USA;

⁴Interdisciplinary Neuroscience Program and MD-PhD Program, Yale University School of Medicine, New Haven, Connecticut 06510, USA;

⁵Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD, 20 815



≪

dx.doi.org/10.17504/protocols.io.bp2l618edvqe/v1

William Hancock-Cerutti

This protocol describes an assay to detect motor deficits in mice by testing their ability to traverse a thin dowel between towers.

HC-Protocol-BalanceBeam.docx

DOI

dx.doi.org/10.17504/protocols.io.bp2l618edvqe/v1

William Hancock-Cerutti, Marianna Leonzino, Pietro De Camilli 2022. Balance beam assay to detect motor phenotypes in mice. **protocols.io** https://dx.doi.org/10.17504/protocols.io.bp2l618edvqe/v1

_____ protocol,

Apr 17, 2022

Apr 18, 2022

60896



1

Citation: William Hancock-Cerutti, Marianna Leonzino, Pietro De Camilli Balance beam assay to detect motor phenotypes in mice https://dx.doi.org/10.17504/protocols.io.bp2l618edvge/v1