



VERSION 3

FEB 01, 2024

OPEN ACCESS



**Protocol Citation:** Nimalka Weerasuriya 2024. Pythium Zoospore Production Soaking Solution. [protocols.io](https://protocols.io/view/pythium-zoospore-production-soaking-solution-c8jazuie) <https://protocols.io/view/pythium-zoospore-production-soaking-solution-c8jazuie> Version created by Nimalka Weerasuriya

**License:** This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** In development  
We are still developing and optimizing this protocol

**Created:** Feb 01, 2024

**Last Modified:** Feb 01, 2024

## Pythium Zoospore Production Soaking Solution V.3

Nimalka Weerasuriya<sup>1</sup>

<sup>1</sup>USDA, OSU Stillwater



Nimalka Weerasuriya  
USDA, Oklahoma State University

### DISCLAIMER

#### DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to [protocols.io](https://protocols.io) is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with [protocols.io](https://protocols.io), can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

### ABSTRACT

Creation and test of soaking solutions to be used for large-scale zoospore production for *Pythium myriotylum*. This is modified from methods in:

Nyochembeng, L. M., Pacumbaba, R. P., & Beyl, C. A. (2002). Calcium Enhanced Zoospore Production of *Pythium myriotylum* in vitro. *Journal of Phytopathology*, 150(7), 396–398.

<https://doi.org/10.1046/J.1439-0434.2002.00759.X>

PROTOCOL integer ID: 94530

## MATERIALS

**Keywords:** pythium, myriotylum, zoospore, soaking solutions, fungi, oospore

## Soaking Solutions:

- Verified Pythium myriotylum culture
- 1.5-2% WA or CMA plates, 90 mm (4 \* replicates)
- RO/DI water
- 1 L beaker
- 4 x 500 mL autoclavable bottles and lids
- Calcium carbonate ( $\text{CaCO}_3$ )
- Sucrose
- Whatman #1 filter paper and funnel
- 1 N KOH – pH adjustment
- 1-5 N HCl – pH adjustment
- pH meter and standards
- Stir bars and stir plate
- Autoclave

## Testing Zoospore Soaking Solutions:

- Haemocytometer
- Microscope 40X and slides
- Counter
- 0.08% Methylene blue

## Preparation

- 1 Have mature colonies of verified *Pythium myriotylum* growing on CMA or 1.5-2% WA 90 mm plates. Colony maturity ~7-14 days, with visible oospores.

## Note

Fungal growth rates may vary with ambient temperature and petri plate sizes (60 mm vs 90 mm), so wait until mycelium covers the petri plate and has visible oospores before proceeding. Oospores will form at the edges of the dish.

## CITATION

Jones, B. L., & Woodard, K. E (1986). A Technique for Evaluating Peanut Germ Plasm for Resistance to *Pythium myriotylum*. Plant Disease, 70(11), 1038–1043.

LINK

<https://doi.org/10.1094/PD-70-1038>

## Soaking Solutions

- 2 Make Soaking Solutions 1, 2, 3, and Control.
  - Prep 4 x 1 L autoclavable bottles for each Soaking Solutions (1-3) and Control.







### CITATION

Nyochembeng, L. M., Pacumbaba, R. P., & Beyl, C. A (2002). Calcium Enhanced Zoospore Production of *Pythium myriotylum* in vitro. *Journal of Phytopathology*, 150(7), 396–398.

LINK

<https://doi.org/10.1046/J.1439-0434.2002.00759.X>

### 2.1 Soaking Solution #1, [M] 0.01 Molarity (m) Ca++ (1 L):



-  1 L RO water at  7 in 1 L beaker
- Add  1 g CaCO<sub>3</sub>
- Filter through Whatman #1 filter paper using a funnel into a 1 L bottle.
- Add a stir bar to the bottle.
- Adjust pH from  8.5 to  10.5 to  7.0 with [M] 5 N HCl and [M] 1 Mass Percent KOH to duplicate the soaking solution in original methods.

#### Note

Normality is a measure of concentration that is equal to the gram equivalent weight of solute per litre of solution. Gram equivalent weight is a measure of the reactive capacity of a molecule.


The original methods use 1 N HCl, but it is sometimes too weak to induce pH changes unless large volumes are used.

### 2.2 Soaking Solution #2, [M] 0.01 Molarity (m) Ca++ + [M] 0.001 Molarity (m) sucrose (500 mL):





- Transfer  500 mL of SS#1 into a clean 500 mL bottle labeled SS #2.
- Add  171 mg sucrose to make [M] 0.001 Molarity (m) sucrose solution.

#### Note

Where **Mass (g) = Concentration (mol/L; M) x Volume (L) x Molecular Weight (g/mol)**  
 Sucrose (g) = 0.001 mol/L sucrose x 0.500 L x 342.3 g/mol  
 Sucrose (g) = 171.15 milligrams

- Check that pH is  7.

### 2.3 Soaking Solution #3, 0.001 Molarity (m) sucrose (500 mL):

- Add  171 mg sucrose to make  0.001 Molarity (m) sucrose solution in  500 mL RO water
- Check that pH is  7.

- Autoclave for 20 minutes on liquid cycle. Remove from autoclave immediately to reduce water evaporation and concentration changes.

#### Protocol



NAME

**Sterilizer (Consolidated)**



CREATED BY



Nimalka Weerasuriya

**PREVIEW**

## Testing Zoospore Production

1d

- Take  30 mL of each Soaking Solution  Soaking Solution Preparation onto surface of *Pythium myriotylum* plate cultures that have actively produced oospores.

- Incubate under light at  Room temperature for  24:00:00 .  
Check for abundant sporangia that will appear after immersion.

1d

- After 1.5 up to 4 h every 30 minutes:



Take  $\text{80 } \mu\text{L}$  soaking liquid from each immersion to a microcentrifuge tube with  $\text{20 } \mu\text{L}$  0.08% methylene blue. Gently invert to stain (or gently vortex) and immobilize zoospores.

#### Note

Methylene blue (0.1% stock):

- Dissolve  $\text{0.1 g}$  methylene blue in  $\text{100 mL}$  dH<sub>2</sub>O

Methylene blue (0.08% working solution):

- Mix  $\text{8 mL}$  MB stock solution with  $\text{92 mL}$  RO water in a labeled dropper bottle. This solution is used for the viability staining.

7 Take  $\text{10 } \mu\text{L}$  of liquid into haemocytometer and examine under 40x to quantify.