



4 ▼

May 31, 2022

## 🌐 2-step PCR mixture and conditions (Barcoded-head primers for seqs pooling) V.4

Yin-Tse Huang<sup>1</sup><sup>1</sup>Kaohsiung Medical University

protocol .



Yin-Tse Huang

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to [protocols.io](https://protocols.io) is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with [protocols.io](https://protocols.io), can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

### PCR mixture and condition (2X SUPERGREEN PCR MASTER MIX)

Yin-Tse Huang 2022. 2-step PCR mixture and conditions (Barcoded-head primers for seqs pooling). **protocols.io**  
<https://protocols.io/view/2-step-pcr-mixture-and-conditions-barcoded-head-pr-b99rr956>  
Yin-Tse Huang



protocol

Herbold CW, Pelikan C, Kuzyk O, Hausmann B, Angel R, Berry D, Loy A. 2015. A flexible and economical barcoding approach for highly multiplexed amplicon sequencing of diverse target genes. Front. Microbiol. [Internet] 6:731. Available from: <http://dx.doi.org/10.3389/fmicb.2015.00731>

---

 protocol ,

May 30, 2022

May 31, 2022

63505

:

**DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK**

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to [protocols.io](https://protocols.io) is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with [protocols.io](https://protocols.io), can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

- 1 Wear glove, clean up the working bench w. 1% bleach




For 1' PCR head-primers

- 2 Prepare 1' PCR master mixutre for **head-primers** (prepare 1.2X of solutions for pipetting error if needed)

PCR mixture for head-primers for each reaction

A	B	C	D
Component	Volume	Volume (1.2X)	Final conc.
Forward Primer (10 µM)	1.6 µl	1.9 µl	1 µM
Reverse Primer (10 µM)	1.6 µl	1.9 µl	1 µM
2X Supergreen PCR Master Mix	7.8 µl	9.4 µl	-
ddH2O	4.1 µl	4.9 µl	-
Total volume	15 µl	18 µl	-

- 3 Mix the 1' PCR master mixture gently by pipetting. Quick spin the tube.

- 4 Transfer  **15 µL** 1' PCR master mixutre in 8-strip PCR tubes.
- 5 Add  **0.6 µL** DNA template in 8-strip PCR tubes, resulting  **15.6 µL** reaction mixture for 1' PCR.
- 6 Mix the reaction mixture gently by tapping the tubes. Quick spin the tubes.
- 7 Carry out PCR using the following condition:

1' PCR condition for **head-primers**

A	B	C	D
Step	Temp	Sec	Cycle
<i>Initial denaturation</i>	95 °C	30-180 (a)	
<i>Denaturation</i>	98 °C	15	20-25 cycles
<i>Annealing</i>	64-68 °C varied (b)	15	
<i>Extension</i>	72 °C	60-180 (c)	
<i>Final extension</i>	72 °C	210	
<i>Preservation</i>	Preservation	4 °C	∞

a. Varied depend on template complexity


b. Annealing varied, **62-65°C** is working based on test on 220530; Refer to 1' PCR primers for annealing temperature

c. 1kb ~ 1min extension; enough time allow full extension of sequence

## 7.1 1' hear-primers used in Huang lab


A	B	C	D
Name	Sequence	Tm°C	CG%
NS1B1ngs_H1	gctatgcgcgagctgccctngttgatytgccagt	71.7	60
ITS4ngs_H1	gctatgcgcgagctgctcctscgcttattgatatgc	69	55.6
LR5_H1	gctatgcgcgagctgctcctgagggaaactcg	70.2	60.6
EF1-526F_H1	gctatgcgcgagctgctgctgtygtyatygghcaygt	71	59.3
EF1-1567R_H1	gctatgcgcgagctgcachgtccraccacccratctt	70.6	56
EF1-2218R_H1	gctatgcgcgagctgcatgacaccracrgcracrgtytg	72.2	60.3
Ben2f_H1	gctatgcgcgagctgctccagactggtcagtgtgtaa	70.5	56.8
Bt2b_H1	gctatgcgcgagctgcaccctcagtgtagtacccttggc	74.5	62.5
T22_H1	gctatgcgcgagctgctctggatgttgttggaatcc	70.3	56.8
RPB2-3bF_H1	gctatgcgcgagctgcggwggwtayttyatyatyaatgg	65.6	48.7
RPB2-7cR_H1	gctatgcgcgagctgccccatrgcttgytrcccat	72.3	59.7
fRPB2-11aR_H1	gctatgcgcgagctgcgcrtggatcttrtcrtsacc	71.7	60.8

## 8 Carry out **electrophoresis** for inspection of DNA products



Gel before markdown

## 9 Markdown wells and upload the pictures to the Lab Google drive



Marked gel picture go to the Lab Google drive

For 2' PCR barcoded-head primers

- 10 Prepare 2' PCR master mixutre for **barcoded-primers** (prepare 1.2X of solutions for pipetting error if needed)

PCR mixture for barcoded-primers for each reaction (**NO PRIMERS at this point!!**)

A	B	C	D
Component	Volume	Volume (1.2X)	Final conc.
2X Supergreen PCR Master Mix	10.75 µL	12.9 µL	-
ddH2O	10.75 µL	12.9 µL	-
Total volume	21.5 µL	25.8 µL	-

- 11 Mix the 2' PCR master mixture gently by pippeting. Quick spin the tube.
- 12 Transfer **21.5 µL** of the 2' PCR master mixture to PCR tubes.
- 13 Add **2.5 µL** pre-mixed barcoded-head primers (Forward + Reverse) to each PCR tube
- 14 Add **1 µL** of 1' PCR product as template, resulting in **25 µL** reaction mixture for 2' PCR.

15 Mix gently by tapping the tubes. Quick spin the tubes.

16 Carry out 2' PCR using the following condition:

2' PCR condition for **barcoded-head primers**

A	B	C	D
Step	Temp	Sec	Cycle
<i>Initial denaturation</i>	98 °C	30	
<i>Denaturation</i>	98 °C	15	10-15 cycles
<i>Annealing</i>	64-68 °C varied (a)	15	
<i>Extension</i>	72 °C	60 (b)	
<i>Final extension</i>	72 °C	210	
<i>Preservation</i>	Preservation	4 °C	∞

a. Annealing varied, **XX-XX C** is working based on test on XXXXXX; Refer 2' PCR primers for annealing temperature

b. 1kb ~ 1min extension; enough time allow full extension of sequence

### 16.1 2' barcoded-head primers used in Huang lab

A	B	C	D
Name	Sequence	Tm°C	CG%
F1-1	aagaaagttgtcgggtgtctttgtggctatgcgcgagctgc	70.3	52.5
F1-2	tcgattccgtttgttagtcgtctgtgctatgcgcgagctgc	70.9	55
F1-3	caggtagaaagaagcagaatcgagctatgcgcgagctgc	70	55
F1-4	ttcggattctatcgtgtttccctagctatgcgcgagctgc	69.2	52.5
F1-5	ctgtccagggtttgtgtaaccttgctatgcgcgagctgc	70.7	55
F1-6	ttctcgcaaaggcagaaagtagtcgctatgcgcgagctgc	71.3	55
F1-7	gtgttaccgtgggaatgaatccttgctatgcgcgagctgc	70.6	55
F1-8	ttcagggaacaaaccaagtacgtgctatgcgcgagctgc	70.2	52.5
R1-1	gattctgattactctattcgccaggctatgcgcgagctgc	68.5	52.5
R1-2	ggaataataaccattgaagtagcacgctatgcgcgagctgc	67.5	50
R1-3	ttgctacggttgacctgcagttagctatgcgcgagctgc	71.4	55
R1-4	aacttgaggatcgtatattcaatgctatgcgcgagctgc	65.2	45
R1-5	gggtccctctactcatttagcatggctatgcgcgagctgc	71.4	57.5
R1-6	cagagctgaccctccagatatttggtatgcgcgagctgc	71.5	57.5
R1-7	atagctgaagcaatctacctatcggtatgcgcgagctgc	69.2	52.5
R1-8	cagagtaagggataggttcggcagctatgcgcgagctgc	71.1	57.5
R1-9	caatcaacgaattagatgtcgggtgctatgcgcgagctgc	69.1	52.5
R1-10	gaccttagtcacatggtagtctaagctatgcgcgagctgc	68	52.5
R1-11	gttcggatgcaatatggttactggctatgcgcgagctgc	70.7	55
R1-12	tagcagaagtcctgtaagacctgctatgcgcgagctgc	70.7	55

17 Carry out **electrophoresis** for inspection of DNA products

18 Markdown wells and upload the pictures to the Lab Google drive