



Oct 25, 2020

DNT Detection In Soil

Zhujun Wei¹¹2020 iGEM NEFU China**1** Works for me dx.doi.org/10.17504/protocols.io.bnxtmfnn

2020 iGEM NEFU China

 Zhujun Wei

DOI

dx.doi.org/10.17504/protocols.io.bnxtmfnn

PROTOCOL CITATION

Zhujun Wei 2020. DNT Detection In Soil. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bnxtmfnn>

LICENSE

This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Oct 25, 2020

LAST MODIFIED

Oct 25, 2020

PROTOCOL INTEGER ID

43731

- 1 The 1% bacteria (PYB1s-yqjF4-luxCDABE) were cultured in LB containing 0.1% streptomycin (50 mg/mL) for 24 h with aeration at 37°C to reach a final density of 1×10^8 cells/mL.
- 2 Preparation of 2% Alginate solution, 1% calcium-chloride solution, experimental soil: dissolve 2g Alginate in 100 mL distilled water (2% w/w) and 1g calciumchloride in 100 mL distilled water (1% w/w). The experiment soil was filtrated prior to use.
- 3 Sterilize the 2% Alginate solution, 1% calcium-chloride solution, (121°C, 15min), and experimental soil(121°C, 1.5h).
- 4 Add 0, 0.25, 0.5and 1mg of DNT powder(0, 250, 500 and 1000mg/kg) to the wells of a 12-well plate. Take 1g of experimental soil and drop it onto the DNT in the 12-well plate on a clean bench.
- 5 The engineered bacteria were added into 2% Alginate solution at a 1:9 (v/v) ratio.

- 6 Drop 200 μ L of 1% (w/w) calcium-chloride solution into the 1mL alginate-bacterial suspension droplets (the volume ratio between the Alginate mixture and calcium chloride solution was 5:1). Leave the mix for 1 minute.
- 7 The mixed Alginate gel was tiled on the soil.