



Nov 20, 2021

🌐 Factors associated with admission to the intensive care unit and mortality in patients with COVID-19, Colombia

Jorge Enrique Machado-Alba¹, Luis Fernando Valladales Restrepo¹,
Manuel E Machado Duque¹, Andres Gaviria Mendoza¹

¹Universidad Tecnológica de Pereira

1



dx.doi.org/10.17504/protocols.io.bud5ns86

 Jorge Machado Alba

Introduction: Coronavirus disease 2019 (COVID-19) has affected millions of people worldwide, and several sociodemographic variables, comorbidities and care variables have been associated with complications and mortality.

Objective: [To identify the factors associated with admission to intensive care units \(ICUs\) and mortality in patients with COVID-19 from 4 clinics in Colombia.](#)

Methods: This was a follow-up study of a cohort of patients diagnosed with COVID-19 between March and August 2020. Sociodemographic, clinical (Charlson comorbidity index and NEWS 2 score) and pharmacological variables were identified. Multivariate analyses were performed to identify variables associated with the risk of admission to the ICU and death ($p < 0.05$).

Results: A total of 780 patients were analyzed, with a median age of 57.0 years; 61.2% were male. On admission, 54.9% were classified as severely ill, 65.3% were diagnosed with acute respiratory distress syndrome, 32.4% were admitted to the ICU, and 26.0% died. The factors associated with a greater likelihood of ICU admission were severe pneumonia (OR: 9.86; 95%CI:5.99-16.23), each 1-point increase in the NEWS 2 score (OR:1.09; 95%CI:1.002-1.19), history of ischemic heart disease (OR:3.24; 95%CI:1.16-9.00), and chronic obstructive pulmonary disease (OR:2.07; 95%CI:1.09-3.90). The risk of dying increased in those older than 65 years (OR:3.08; 95%CI:1.66-5.71), in patients with acute renal failure (OR:6.96; 95%CI:4.41-11.78), admitted to the ICU (OR:6.31; 95%CI:3.63-10.95), and for each 1-point increase in the Charlson comorbidity index (OR:1.16; 95%CI:1.002-1.35).

Conclusions: Factors related to increasing the probability of requiring ICU care or dying in patients with COVID-19 were identified, facilitating the development of anticipatory intervention measures that favor comprehensive care and improve patient prognosis.

[BD_Morbimortalidad-COVID-19.xlsx](#)

DOI

dx.doi.org/10.17504/protocols.io.bud5ns86

Jorge Enrique Machado-Alba, Luis Fernando Valladales Restrepo, Manuel E Machado Duque, Andres Gaviria Mendoza 2021. Factors associated with admission to the intensive care unit and mortality in patients with COVID-19, Colombia. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bud5ns86>



protocol

Machado-Alba JE, Valladales-Restrepo LF, Machado-Duque ME, Gaviria-Mendoza A, Sánchez-Ramírez N, Usma-Valencia AF, Rodríguez-Martínez E, Rengifo-Franco E, Forero-Supelano VH, Gómez-Ramírez DM, Sabogal-Ortiz A (2021) Factors associated with admission to the intensive care unit and mortality in patients with COVID-19, Colombia. PLoS ONE 16(11): e0260169. doi: [10.1371/journal.pone.0260169](https://doi.org/10.1371/journal.pone.0260169)

protocol ,

Apr 21, 2021

Nov 20, 2021

49309

1