



APR 10, 2024

Morphometry of thyroid cartilage, epiglottis and pyriform fossa

Murlimanju Virupakshamurthy¹

¹Department of Anatomy, Kasturba Medical College, Mangalore, Manipal Academy of Higher Education, Manipal, Karnataka, India.



Murlimanju Virupakshamurthy

Department of Anatomy, Kasturba Medical College, Mangalore, ...

ABSTRACT

Larynx is a complex organ of voice production and respiration, which is supported by a series of cartilages, membranes, muscles and joints. They bring about movement of vocal cords with a considerable range of mobility. Apart from the phonating mechanism, it also serves as a sphincter or watch dog of the lower respiratory tract. The knowledge of laryngeal anatomy is important to various professionals such as phoniaticians, speech therapists, oncologists and otorhinolaryngologists. Understanding the laryngeal framework is mandatory in constructing biomechanical models for voice disorders that requires accurate laryngeal dimensions. The data of the dimensions of thyroid cartilage, epiglottis and piriform fossa are essential to the anesthesiologists during the laryngoscopy and endotracheal intubation. However, they are scarce in the scientific literature. There are not many studies available about the measurements of thyroid cartilage, piriform fossa and epiglottis in our sample population.

ATTACHMENTS

[larynx protocols.docx](#)

OPEN ACCESS



DOI:

dx.doi.org/10.17504/protocols.io.x54v92m8ml3e/v1

Protocol Citation: Murlimanju Virupakshamurthy 2024. Morphometry of thyroid cartilage, epiglottis and pyriform fossa.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.x54v92m8ml3e/v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: Apr 09, 2024

Last Modified: Apr 10, 2024

PROTOCOL integer ID: 97946

Keywords: Piriform Fossa,
Thyroid Cartilage, Epiglottis

PROTOCOL REFERENCES

Ajmani

ML, Jain SP, Saxena SK: A metrical study of laryngeal cartilages and their ossification. *Anat. Anz.* 1980; 148(1): 42-48.

Anson

BJ, McVay CB: *Surgical Anatomy*. 5th ed. WB Saunders; 1971; 255.

Chau

CW, Chan HH, Wong CP, Wong TW, Lau CC: Sonographic measurement of the epiglottis in normal Chinese adults. *Hong Kong J. Emerg. Med.* 2010; 17(5): 429-434.

Davis EDD: The applied anatomy and

physiology of the pharynx and oesophagus. *Ann. R. Coll. Surg. Engl.* 1948; 3(3): 139-153.

Eckel HE, Sittel C, Zorowka P, Jerke A:

Dimensions of laryngeal framework in adults. *Surg. Radiol. Anat.* 1994; 16(1): 31-36.

Hajjiioannou JK, Florou V, Kousoulis P: Superior

thyroid cornu anatomical variation causing globus pharyngeous and dysphagia. *Case Rep. Med.* 2010; 2010: 142928.

Jain

M, Dhall U: Morphometry of the thyroid and cricoid cartilages in adults on C.T. scan. *J. Anat. Soc. Ind.* 2010; 59(1): 19-23.

Lang J, Fischer K, Nachbaur S: Uber Masse, Form

und Formvarianten der Cartilagine thyreoidea et cricoidea [Measurements, form and form variants of the thyroid and cricoid cartilages]. *Gegenbaurs. Morphol. Jahrb.* 1984; 130(5) :639-657.

Park

DK, Kim DI, Lee UY, Han KH, Kim KH, Han SH: Morphometric analysis of the Korean thyroid cartilage for identification of sex: metric study. *Korean J. Phys. Anthropol.* 2003; 16(3): 129-136.

Randestad

A, Lindholm CE, Fabian P: Dimensions of the cricoid cartilage and the trachea. *Laryngoscope.* 2000; 110(11): 1957-1961.

Savkovic A, Delic J, Isakovic E, Ljuca F, Nuhbegovic S: Age

characteristics of the larynx in infants during the first year of life. *Periodicum Biologorum* 2010; 112(1): 75-82.

Werner

SL, Jones RA, Emerman CL. Sonographic assessment of the epiglottis. *Acad. Emerg. Med.* 2004; 11(12): 1358-1360.

Zrunek

M, Happak W, Hermann M, Strienzer W: Comparative anatomy of human and sheep laryngeal skeleton. Acta Otolaryngol. 1988; 105(1-2): 155-162.

GUIDELINES

This study was approved by the ethics committee of our institution (Approval Committee Name: Institutional Ethics Committee, Kasturba Medical College, Mangalore, Approval Number: IEC KMC MLR: 02/2022/61, dated 17.02.2022)

MATERIALS

Study setting: Department of Anatomy, Kasturba Medical College, Mangalore

Study participants:

Human embalmed cadavers

Inclusion criteria: Adult human embalmed cadavers

Study duration: 3 months

Sample size: 22

Sampling method: The sample size is similar to the earlier study performed by Eckel et al. (1994).

Data collection methodology:

A digital Vernier caliper (Mitutoyo, Japan, 0-150 mm 500-196) will be used to perform the measurements. Height, width and thickness of the thyroid cartilage, epiglottis and piriform fossa will be measured. The measurements will be taken at their maximum diameter and for the thyroid cartilage and piriform fossa, the readings will be taken separately over the right and left sides.

Data analysis: The

data will be presented as mean \pm standard deviation and the side-based comparison of the right and left sides will be performed by using the paired 't' test. The recent version of SPSS (version 27) software will be utilized for the statistical analysis. The gender-based comparison and age wise segregation will not be performed in the present study.

Morphometry of thyroid cartilage, epiglottis and pyriform fossa

- 1 The goal of this study will be to obtain the dimensions of thyroid cartilage, epiglottis and piriform sinus in embalmed cadavers of Indian sample population.