



VERSION 3

APR 07, 2023

OPEN ACCESS

Protocol Citation: Lynn Doran, Amanda P. De Souza 2023. Total Starch Enzymatic Digestion . [protocols.io](https://protocols.io/view/total-starch-enzymatic-digestion-csbkwakw) <https://protocols.io/view/total-starch-enzymatic-digestion-csbkwakw> Version created by Lynn Doran

MANUSCRIPT CITATION: Amaral, LV do, Gaspar, M., Costa, PMF, Aidar, MPM, & Buckeridge, MS (2007). New fast and sensitive enzymatic method for extracting and measuring starch in plant materials. *Hoeheia*, 34 n.4, 425-431. doi:10.1590/s2236-89062007000400001

Pak S. Chow, Simon M. Landhäusser, A method for routine measurements of total sugar and starch content in woody plant tissues, *Tree Physiology*, Volume 24, Issue 10, October 2004, Pages 1129-1136, <https://doi.org/10.1093/treephys/24.10.1129>


License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

🌐 Total Starch Enzymatic Digestion V.3

Lynn Doran¹, Amanda P. De Souza¹

¹Realizing Increased Photosynthetic Efficiency (RIPE)

UIUC Long Lab



Lynn Doran

Realizing Increased Photosynthetic Efficiency (RIPE)

ABSTRACT

Enzymatic digestion of total soluble starch to glucose in plant tissue extracts for preparation for quantification via the GOD-POD Method (NZYtech).

MATERIALS

Reagents

- [α-amylase \(Bacillus licheniformis\) enzyme, 3000 U/mL](#)
- [Amyloglucosidase \(Aspergillus niger\) enzyme, 3260 U/mL](#)
- MOPS Buffer 10mM, pH 6.5

2.09 g of MOPS per 1 L of water (0.4598 g for 220 mL water), pH adjusted to 6.5 using NaOH.

Store refrigerated for up to 2 months.

- Acetate Buffer 100mM , pH 4.5

6 mL of acetic acid per 1 L of water (1.5 mL per 250 mL water), pH adjusted to 6.5 using NaOH.

Store refrigerated for up to 2 months.

- Ice

Materials

- Pipette tips

Equipment

- Graduated cylinder
- Water bath
- Floating tube holder
- Single channel pipette
- Ice bucket

Protocol status: Working
We use this protocol and it's working

BEFORE START INSTRUCTIONS

Extract and dry total starch pellet from plant tissue per [Extraction of Non-Structural Carbohydrates \(Total Soluble Sugars + Starch\) in Plant Tissues](#).

Created: Apr 03, 2023

Last Modified: Apr 07, 2023

PROTOCOL integer ID:
79948

Keywords: Starch, Soluble
Starch, Enzymatic Digestion,
Plant Tissue, GOD-POD

- 1 Prepare fresh daily 120 U/mL α -amylase in MOPS buffer. 1 mL per sample will be needed. Initial concentration of α -amylase is 3000 U/mL. Use $C_1V_1 = C_2V_2$ to calculate the volume of α -amylase and MOPS buffer to use.

Note


α -amylase comes at 3000 U/mL concentration in a 20 mL bottle. If using all of the bottle, 20 mL of α -amylase in 146.7 mL of MOPS will result in 120 U/mL α -amylase in MOPS buffer.

To prepare 250 mL of 120 U/mL α -amylase in MOPS buffer, dilute 10 mL α -amylase (3000 U/mL) into 240 mL MOPS buffer.

- 2 Prepare fresh daily 30 U/mL amyloglucosidase in acetate buffer. 1 mL per sample will be needed. Initial concentration of amyloglucosidase is 3260 U/mL. Use $C_1V_1 = C_2V_2$ to calculate the volume of amyloglucosidase and acetate buffer to use.

Note

To prepare 250 mL of 30 U/mL amyloglucosidase in acetate buffer, dilute 2.3 mL amyloglucosidase (3260 U/mL) into 247.7 mL acetate buffer.


- 3 Heat the water bath to  75 °C .

- 4 Add  500 μ L of α -amylase in MOPS buffer (120 U/mL) to each sample tube.

5 Vortex to suspend all solids. Flicking the tube may help dislodge the dried starch pellet.

6 Incubate for  00:30:00 at  75 °C in the water bath.

30m

7 Add another  500 μ L of α -amylase in MOPS buffer (120 U/mL) to each sample tube.

8 Vortex to suspend all solids.


9 Incubate for  00:30:00 at  75 °C in the water bath.

30m






10 Cool down the water bath to  50 °C, tubes can be stored at room temp on the counter while water bath cools.

Note

Leaving the lid off, especially if it is a shaking water bath, will help cool. For faster cooling, remove some water from the water bath and refill with cool distilled water. Do not use tap water as mineral buildup or heated chlorine could damage the water bath.

11 Add  500 μ L amyloglucosidase in acetate buffer (30 U/mL) to each sample tube.

12 Vortex to suspend all solids.

- 13** Incubate for  00:30:00 at  50 °C in the water bath. 30m
- 14** Add another  500 µL amyloglucosidase in acetate buffer (30 U/mL) to each sample tube.
- 15** Vortex to suspend all solids.
- 16** Incubate for  00:30:00 at  50 °C in the water bath. 30m
- 17** Place the tubes in ice to stop the reaction, until cool to the touch.
- 18** Proceed to total starch (as glucose) quantification by NZYtech GOD-POD method or store the samples at -20°C up to one month.