



Oct 24, 2020

Specificity test of toehold

Hung Liang Pai¹, Cheng-Ruei Yang¹¹Chung Shan Medical University

1

Works for me

This protocol is published without a DOI.

1

Cheng-Ruei Yang

PROTOCOL CITATION

Hung Liang Pai, Cheng-Ruei Yang 2020. Specificity test of toehold. **protocols.io**
<https://protocols.io/view/specificity-test-of-toehold-bntxmepn>

LICENSE

This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Oct 23, 2020

LAST MODIFIED

Oct 24, 2020

PROTOCOL INTEGER ID

43607

Preparation

- 1 Sterilize the bench, and put on a labmat
- 2 Thaw the reagents including 1. DNase/ RNase free water (store in -20°C) 2. solution A (store in -80°C) 3. solution B (store in -80°C) 4. RNase inhibitor (store in -20°C) 5.toehold switches with invertase DNA (store in 4°C) 6.miRNA as the trigger (store in 4°C) [On ice](#)

Protocols

2h 30m 20s

- 3 Gently shake and spin down the tube before adding (especially solution A and DNA plasmid)
- 4 Add below reagents in to PCR tubes in order:
 1. ddH₂O(till totally [5 µl](#))
 2. solution A [2 µl](#)
 3. solution B [1.5 µl](#)
 4. inhibitor [0.2 µl](#)
 5. toehold switches with invertase DNA [50 ng](#)
 6. miRNA [0 µl](#) [50 µl](#) [100 µl](#) [150 µl](#) [200 µl](#)

- 5 Centrifuge [00:00:20](#) [4000 rpm](#)

20s

- 6 Place the PCR tubes in the dry bath incubator 🔧 **37 °C** ⌚ **02:00:00** 2h
- 7 Put the PCR tubes 🔧 **On ice** after 2 hours
- 8 Measure the glucose concentrations of each PCR tubes by glucometer
- 9 Put the PCR tubes in the dry bath incubator 🔧 **55 °C** ⌚ **00:30:00** 30m
- 10 Measure the glucose concentrations of each PCR tubes by glucometer again