



Apr 09, 2021

Genome-wide quantification of TF binding at single DNA molecule resolution

Rozemarijn Kleinendorst¹, [guido.barzaghi](#)^{1,2}, Mike L. Smith¹, Judith B. Zaugg³, Arnaud Krebs¹¹European Molecular Biology Laboratory (EMBL), Genome Biology Unit, Meyerhofstraße 1, 69117 Heidelberg;²Faculty of Biosciences, Collaboration for Joint PhD Degree between EMBL and Heidelberg University, Heidelberg, Germany;³European Molecular Biology Laboratory (EMBL), Structural and Computational Biology Unit, Meyerhofstraße 1, 69117 Heidelberg

Rozemarijn Kleinendorst: These authors contributed equally to this work;

[guido.barzaghi](#) : These authors contributed equally to this work;Arnaud Krebs: Correspondence should be addressed to: arnaud.krebs@embl.de**1** Works for medx.doi.org/10.17504/protocols.io.btr2nm8e[Krebs_EMBL](#)[guido.barzaghi](#)

ABSTRACT

Precise control of gene expression requires the coordinated action of multiple factors at cis-regulatory elements (CREs). We recently developed Single Molecule Footprinting (SMF) to simultaneously resolve the occupancy of multiple proteins including Transcription Factors (TFs), RNA Pol II (Pol II) and nucleosomes on single DNA molecules genome-wide. The technique combines the use of cytosine methyltransferases to footprint the genome with bisulfite sequencing to resolve TF binding patterns at CREs. DNA footprinting is performed by incubating permeabilized nuclei with recombinant methyltransferases. Upon DNA extraction, whole genome or targeted bisulfite libraries are prepared and loaded on Illumina sequencers. The protocol can be completed in 4-5 days. Analysis can be performed in 2 days using a dedicated R package. The protocol can be executed in any laboratory with access to high-throughput sequencing. Our method can be used to analyse how TFs cooperate and antagonize to regulate transcription.

ATTACHMENTS

[Kleinendorst_et_al_preprint.pdf](#)

DOI

dx.doi.org/10.17504/protocols.io.btr2nm8e

PROTOCOL CITATION

Rozemarijn Kleinendorst, [guido.barzaghi](#), Mike L. Smith, Judith B. Zaugg, Arnaud Krebs 2021. Genome-wide quantification of TF binding at single DNA molecule resolution. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.btr2nm8e>

KEYWORDS

chromatin accessibility, DNA footprinting, Transcription Factor, gene regulation, RNA Pol II, DNA methylation.

LICENSE

———— This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Mar 29, 2021

LAST MODIFIED

Apr 09, 2021

PROTOCOL INTEGER ID

48666