



SEP 19, 2023

🌐 Double-fixation prior to ChIPs

Jacob G

Kirkland¹,

Mary Bergwell¹, JinYoung Park^{1,2}

¹Oklahoma Medical Research Foundation;

²University of Oklahoma Health Sciences Center



Jacob G Kirkland

Oklahoma Medical Research Foundation, Oklahoma University He...

DISCLAIMER

OPEN  ACCESS



DOI:

dx.doi.org/10.17504/protocols.io.yxmvm396nl3p/v1

Protocol Citation: Jacob G Kirkland, Mary Bergwell, JinYoung Park 2023. Double-fixation prior to ChIPs.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.yxmvm396nl3p/v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Sep 19, 2023

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to [protocols.io](#) is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with [protocols.io](#), can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

ABSTRACT

This protocol is for the double-fixation of chromatin regulators for chromatin precipitation. The double-fixation method can improve signal-to-noise ratios of chromatin regulator ChIPs, especially for transiently bound proteins or those found in large protein complexes.

MATERIALS

Disuccinimidyl glutarate (DSG) [ChemCruz]

DMSO [Sigma]

PBS [Fisher Scientific]

2.5M Glycine Solution: Dissolve 93.84g Glycine (MW: 75.07g/mol) in 400ml sterile water. Adjust final volume to 500 ml and sterile filter

CiA Fix Buffer: 25 ml 1M HEPES pH 8.0; 1 ml 0.5M EDTA; 0.5 ml 0.5M EGTA; 10 ml 5M NaCl Adjust final volume to 500 ml with sterile water and filter sterilize

trypsin-EDTA 0.25% (Fisher Scientific)

16% Formaldehyde Solution (w/v) Methanol-Free [Thermo Scientific]

SAFETY WARNINGS



Use proper safety precautions when using formaldehyde.

Fresh DSG

30m

- 1 Place DSG at room temperature for approximately 30 minutes before weighing it out.

30m

- 2 Measure DSG and dissolve in DMSO accordingly (using approximately 80 ul of solution per sample)

5m

	Number of Samples	DSG (mg)	Vol. DMSO (ul)
	2	13	160
	4	25	320
	6	38	480
	12	72	960

Isolate and count single cells

20m

- 3 Harvest the cells by first washing with PBS 1m
- 4 Incubate cells in trypsin-EDTA 0.25% until dissociated 5m
- 5 Quench trypsin with media + FBS and move cells to a 15ml conical. Spin down at 300g for 4 minutes 4m
- 6 Resuspend cells in PBS and count 5m
- 7 Move 30e6 cells to a new tube and bring the volume up to 10mls with PBS and spin down at 300g for 4 minutes 4m
- 8 Resuspend cells in 10mls fresh PBS 1m

DSG fixation (First)

36m

- 9 Add 80 ul DSG stock to each sample in 10 ml of PBS and incubate for 30 minutes at room temperature with rocking. 30m


Note: After incubation with DSG, cells may adhere to the walls of pipette tips, so following this incubation, it is recommended that tips be coated with 1% BSA/PBS

- 10 Pellet samples by spinning at 1500 x g for 10 minutes at 4C 4m
- 11 Carefully aspirate the supernatant, taking care not to disturb the pellet. 1m
- 12 Resuspend each sample in 10 ml of CiA Fix Buffer 1m

Formaldehyde Fixation (Second)

34m

- 13 Add 667 ul of 16% Formaldehyde to the sample dropwise and incubate for exactly 10 minutes at room temperature using rotation 10m
- 14 Stop cross-linking by adding 555 ul of 2.5M glycine. 1m
- 15 Incubate samples on ice for 5 minutes 5m
- 16 Pellet samples by spinning at 1500 x g for 10 minutes at 4°C 10m
- 17 Aspirate supernatant taking care not to disturb the pellet 1m

- 
- 18** Wash fixed cells with 10ml PBS 1m
 - 19** Pellet samples by spinning at 1000 x g for 5 minutes at 4C and aspirate supernatant. 5m
 - 20** Samples can be stored at -80°C (snap freeze) or can begin processing for chromatin immunoprecipitation 1m