



Sep 06, 2022

Detection of Tau ubiquitylation

Itika Saha¹, F. Ulrich Hartl^{2,3}, Mark S. Hipp^{2,3}¹Department of Cellular Biochemistry, Max Planck Institute of Biochemistry, Am Klopferspitz 18, 82152 Martinsried, Germany;²Department of Biomedical Sciences of Cells and Systems, University Medical Center Groningen, University of Groningen, Antonius Deusinglaan, 1, 9713 AV Groningen, The Netherlands;³School of Medicine and Health Sciences, Carl von Ossietzky University Oldenburg, 26129 Oldenburg, Germany

1 Works for me

Share

dx.doi.org/10.17504/protocols.io.3byl4je4rlo5/v1

Felix Kraus

ABSTRACT

This protocol describes the detection of ubiquitylated Tau from HEK293 cells stably expressing and propagating aggregates of Tau repeat domain fused to YFP (Sanders et al. Neuron, 2014; Saha et al, BioRxiv, 2022).

DOI

dx.doi.org/10.17504/protocols.io.3byl4je4rlo5/v1

EXTERNAL LINK

<https://www.biorxiv.org/content/10.1101/2022.02.18.481043v1.full>

PROTOCOL CITATION

Itika Saha, F. Ulrich Hartl, Mark S. Hipp 2022. Detection of Tau ubiquitylation.
protocols.io
<https://protocols.io/view/detection-of-tau-ubiquitylation-cf7vtrn6>

MANUSCRIPT CITATION please remember to cite the following publication along with this protocol

The AAA+ chaperone VCP disaggregates Tau fibrils and generates aggregate seeds Itika Saha, Patricia Yuste-Checa, Miguel Da Silva Padilha, Qiang Guo, Roman Körner, Hauke Holthausen, Victoria A. Trinkaus, Irina Dudanova, Rubén Fernández-Busnadiego, Wolfgang Baumeister, David W. Sanders, Saurabh Gautam, Marc I. Diamond, F. Ulrich Hartl, Mark S. Hipp bioRxiv 2022.02.18.481043; doi: <https://doi.org/10.1101/2022.02.18.481043>

LICENSE

————— This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Sep 05, 2022

LAST MODIFIED

Sep 06, 2022

PROTOCOL INTEGER ID

69589

- 1 Harvest cells from 2 confluent wells of a 6 well plate.
- 2 Lyse aggregate-containing cells by vortexing in cold RIPA buffer (Thermo) supplemented with protease inhibitor cocktail (Roche) and DNase. 20 mM N-ethylmaleimide should be included in the lysis buffer to inhibit the activity of deubiquitinating enzymes.
- 3 Briefly sonicate lysate and centrifuge at 2,000 x g for 5 min to remove cell debris. 5m
- 4 Collect supernatant, determine protein concentration and normalize across samples.
- 5 Dilute 1 mg protein in a total volume of 600 µL RIPA buffer.
- 6 Add 50 µL anti-GFP bead slurry (µMACS GFP Isolation kit, Miltenyi Biotec) to diluted lysate.
- 7 Incubate for 1 h at 4 °C in a rotating wheel at 10 rpm. 1h

- 8 Before the end of 1 h, place μ -columns (Miltenyi Biotec) in the magnetic field of a μ MACS Separator (Miltenyi Biotec) and equilibrate columns by applying 250 μ L RIPA buffer. Allow complete flow-through.
- 9 Apply cell lysates and beads to μ -columns. Allow complete flow-through.
- 10 Wash columns 4 times with 1 mL 0.1% SDS/PBS.
- 11 Elute by applying 50 μ L pre-heated (95 °C) 1x SDS sample buffer.
- 12 Analyze eluates by immunoblotting with antibodies against GFP or ubiquitin.