A



Jan 05, 2021

# Protocols for "Shotgun Metagenomics of 361 elderly women reveals gut microbiome change in bone mass loss"

Forked from Protocols for "Shotgun Metagenomics of 361 elderly women reveals gut microbiome change in bone mass loss"

Qi Wang<sup>1,2</sup>, Qiang Sun<sup>1,3</sup>, Xiaoping Li<sup>1</sup>, Zhefeng Wang<sup>4</sup>, Haotian Zheng<sup>1,2</sup>, Yanmei Ju<sup>1,2</sup>, Ruijin Guo<sup>1,5</sup>, Songlin Peng<sup>4</sup>, Huijue Jia<sup>1,6</sup>

<sup>1</sup>BGI-Shenzhen, Shenzhen 518083, China;

<sup>2</sup>School of Future Technology, University of Chinese Academy of Sciences, Beijing, 101408, China.;

<sup>3</sup>Department of Statistical Sciences, University of Toronto, Toronto, Canada;

<sup>4</sup>Department of Spine Surgery, Shenzhen People's Hospital, Ji Nan University Second College of Medicine, 518020, Shenzhen, Chin a.:

<sup>5</sup>Macau University of Science and Technology, Taipa, Macau 999078, China;

<sup>6</sup>Shenzhen Key Laboratory of Human Commensal Microorganisms and Health Research, BGI-Shenzhen, Shenzhen 518083, China

1 Works for me dx.doi.org/10.17504/protocols.io.bq6hmzb6

BGI GIGA 1 more workspace

wangqi

#### **ABSTRACT**

Bone mass loss contributes to the risk of bone fracture in the elderly. Many factors including age, obesity, estrogen and diet, are associated with bone mass loss. Mice studies suggested that the gut microbiome might affect the bone mass by regulating the immune system, however there has been little evidence from human studies. Bone loss increases after menopause. Therefore, we have recruited 361 Chinese post-menopausal women to collect their fecal samples and metadata to conduct metagenome-wide association study (MWAS) to investigate the influence of the gut microbiome on bone health. Gut microbiome sequencing data were produced using BGISEQ500 sequencing, Bone mineral density (BMD) was calculated using Hologic dual energy X-ray machine, body mass index (BMI) and age were also recorded. This collected data allows exploration of the gut microbial diversity and their links to bone mass loss, as well as microbial markers for bone mineral density. In addition, these data are potentially useful in studying the role the gut microbiota might play in bone mass loss and in exploring the bone mass loss process.

DOI

dx.doi.org/10.17504/protocols.io.bq6hmzb6

**COLLECTION CITATION** 

Qi Wang, Qiang Sun, Xiaoping Li, Zhefeng Wang, Haotian Zheng, Yanmei Ju, Ruijin Guo, Songlin Peng, Huijue Jia 2021. Protocols for "Shotgun Metagenomics of 361 elderly women reveals gut microbiome change in bone mass loss". **protocols.io** 

https://dx.doi.org/10.17504/protocols.io.bq6hmzb6

FORK NOTE

FORK FROM

Forked from Protocols for "Shotgun Metagenomics of 361 elderly women reveals gut microbiome change in bone mass loss", wangqi

#### LICENSE

This is an open access collection distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Jan 05, 2021

LAST MODIFIED

Jan 05, 2021

# OWNERSHIP HISTORY

### COLLECTION INTEGER ID

45993

# ABSTRACT

Bone mass loss contributes to the risk of bone fracture in the elderly. Many factors including age, obesity, estrogen and diet, are associated with bone mass loss. Mice studies suggested that the gut microbiome might affect the bone mass by regulating the immune system, however there has been little evidence from human studies. Bone loss increases after menopause. Therefore, we have recruited 361 Chinese post-menopausal women to collect their fecal samples and metadata to conduct metagenome-wide association study (MWAS) to investigate the influence of the gut microbiome on bone health. Gut microbiome sequencing data were produced using BGISEQ500 sequencing, Bone mineral density (BMD) was calculated using Hologic dual energy X-ray machine, body mass index (BMI) and age were also recorded. This collected data allows exploration of the gut microbial diversity and their links to bone mass loss, as well as microbial markers for bone mineral density. In addition, these data are potentially useful in studying the role the gut microbiota might play in bone mass loss and in exploring the bone mass loss process.

