



Mar 25, 2021

Crop FVC retrieval

Liang Wan¹, Jiangpeng Zhu¹, Xiaoyue Du¹, Jiafei Zhang¹, Xiongze Han², Weijun Zhou³, Xiaopeng Li⁴, Jianli Liu⁴, Fei Liang⁵, Yong He¹, Haiyan Cen¹

¹College of Biosystems Engineering and Food Science, and State Key Laboratory of Modern Optical Instrumentation, Zhejiang University, Hangzhou 310058, P.R. China;

²Department of Biosystems Engineering, College of Agriculture and Life Sciences, Kangwon National University, Chuncheon, Kangwon 24341, Korea;

³College of Agriculture and Biotechnology, Zhejiang University, Hangzhou 310058, P.R. China;

⁴Institute of Soil Science, Chinese Academy of Sciences, Nanjing 210008, P. R. China;

⁵Institute of Farmland Water Conservancy and Soil-fertilizer, Xinjiang Academy of Agricultural and Reclamation Science, Shihezi 832000, P.R. China

Liang Wan: 1;

Jiangpeng Zhu: 1;

Xiaoyue Du: 1;

Jiafei Zhang: 1;

Xiongze Han: 2

Weijun Zhou: 3

Xiaopeng Li: 4

Jianli Liu: 4

Fei Liang: 5

Yong He: 1

Haiyan Cen: 1,*

1

Works for medx.doi.org/10.17504/protocols.io.btmynk7w Liang Wan

SUBMIT TO PLOS ONE

ABSTRACT

This protocol describes the method of crop fractional vegetation cover (FVC) retrieval from the PROSAIL-GP model. This method can be applied to UAV-based multispectral images to accurately estimate crop FVC in oilseed rape, rice, cotton and wheat.

DOI

dx.doi.org/10.17504/protocols.io.btmynk7w

PROTOCOL CITATION

Liang Wan, Jiangpeng Zhu, Xiaoyue Du, Jiafei Zhang, Xiongze Han, Weijun Zhou, Xiaopeng Li, Jianli Liu, Fei Liang, Yong He, Haiyan Cen 2021. Crop FVC retrieval. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.btmynk7w>

KEYWORDS

FVC, PROSAIL-GP, crop, UAV, Multispectral images

LICENSE

This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Mar 25, 2021

LAST MODIFIED

Mar 25, 2021

PROTOCOL INTEGER ID

48536

GUIDELINES

Here, the canopy reflectance can be input, which can match with the simulated reflectance. Canopy reflectance is determined based on a weighted combination of the bi-directional and hemispherical-directional reflectance with weights corresponding to the fraction of diffuse incident solar radiation (skyl). The detailed information can be found in chi2P5B.m and PRO4SAIL.m.

In addition, the retrieval results may be slightly affected by the iterative optimization function, while they don't produce the significant difference for model inversion.

The detailed descriptions about the datasets can be found in our published papers.

MATERIALS TEXT


The code of the PROSAIL-GP model and all of the datasets are available on <https://github.com/WanLiangZJU/Crop-FVC-retrieval>.

BEFORE STARTING

The source code of the PROSAIL model used in this study, are open and available on <http://teledetection.ipgp.jussieu.fr/prosail/>. The software for the PROSAIL inversion is open in ARTMO (<https://artmotoolbox.com/>).

1 Make sure you have MATLAB 2014b (or newer) installed on your computer.

2 Download the MATLAB scripts from <https://github.com/WanLiangZJU/Crop-FVC-retrieval>



WanLiangZJU Update README.md 5bb0ada · 5 days ago · 37 commits

File	Action	Time
Dataset	Add files via upload	5 days ago
PROSAIL_5B_GP	Add files via upload	6 days ago
README.md	Update README.md	5 days ago

README.md

Crop-FVC-retrieval

The source code of the PROSAIL model used in this study, are open and available on <http://teledetection.ipgp.jussieu.fr/prosail/>. The software for the PROSAIL inversion is open in ARTMO (<https://artmotoolbox.com/>).

Implementation of model inversion

Please use the code in the file of PROSAIL_5B_GP. Here, we provide five main codes for different datasets as follows: Main_oilseed_rape_2017_2018.m Main_oilseed_rape_2019.m Main_rice_2018.m Main_wheat_2019.m Main_cotton_2019.m

Please update the code and run the main code

1. update the file for the input of reflectance data
2. set the range of model parameters
3. obtain the retrieval result, namely FVC, related to different leaf angle distributions As for different datasets, it is critical to set the suitable range of mode parameters based on previous studies and field measurements.

Please pay attention to some key information This code uses the UAV-based multispectral image with the spectral region of 604-872 nm. If you use different spectral regions, please refer to the original code online <http://teledetection.ipgp.jussieu.fr/prosail/> It is necessary to modify the range of spectral reflectance in the model to fit with your data.

Here, the canopy reflectance can be input, which can match with the simulated reflectance. Canopy reflectance is determined based on a weighted combination of the bi-directional and hemispherical-directional reflectance with weights corresponding to the fraction of diffuse incident solar radiation (skyl). The detailed information can be found in

- 3 Please use the code in the file of **PROSAIL_5B_GP**. Here, we provide five main codes for different datasets as follows: **Main_oilseed_rape_2017_2018.m**, **Main_oilseed_rape_2019.m**, **Main_rice_2018.m**, **Main_wheat_2019.m**, **Main_cotton_2019.m**



- 4 Please update the code and run the main code
1. update the file for the input of reflectance data
 2. set the range of model parameters
 3. obtain the retrieval result, namely FVC, related to different leaf angle distributions. As for different datasets, it is critical to set the suitable range of model parameters based on previous studies and field measurements.

```
clear all; clear all; clc;

for p=1:10 % number of sample
    %% input canopy reflectance (name of excel file)
    %% Notably, as for the canopy reflectance with different bands, we must modify the original data in dataSpect_P5B
    %% dataSpect_P5B includes some prior information about leaf and canopy parameters in the region of 400-2500 nm
    %% to fit with different range of spectral bands, this file should be changed
    R=xlsread('F:\2018水稻\images\biomass-rice2018\FVC+LAI+PROSAIL\JEB\public\Dataset\oilseed rape 2017-2018 dataset.xlsx','spectra');
    R1=R(p,1:25);
    rmes=R1';

    %% determine the LAD
    %% LAD=1, Planophile; LAD=2, Erectophile; LAD=3, Plagiophile;
    %% LAD=4, Extremophile; LAD=5, Spherical; LAD=6, Uniform;
    LAD=6;

    %% The initial value for leaf parameters, which doesn't significantly affect the retrieval.
    Cab=40; % chlorophyll content (ug.cm-2)
    Car=6; % carotenoid content (ug.cm-2)
    Cbrown=0.0; % brown pigment content (ug.cm-2)
    Cw=0.015; % EVI ((g.cm-2)
    Cm=0.004; % LMA (g.cm-2)
    N=1.5; % leaf structure parameter

    %% The initial value for canopy parameters, which doesn't significantly affect the retrieval.
    LAI =3.; % leaf area index (m^2/m^2)
    tts=30; % solar zenith angle (sza)
    tto=0; % viewing zenith angle (vza)
    psi=150; % relative azimuth angle(raa?)
    psoil=0; % soil factor (psoil=0: wet soil / psoil=1: dry soil)
    hspot =0.25;% hot spot
    %% set the range of model parameters based on previous studies and field measurements
```

- 5 This code uses the UAV-based multispectral image with the spectral region of 604-872 nm. If you use different spectral regions, please refer to the original code online <http://teledetection.ipgp.jussieu.fr/prosail/>. It is necessary to modify the range of spectral reflectance in the model to fit with your data.

