

Aug 29, 2024

Kinase activity assays Src and CK2

DOI

dx.doi.org/10.17504/protocols.io.5jyl82by7l2w/v1

Elias Adriaenssens¹

¹Sascha Martens lab, University of Vienna, Max Perutz Labs - Vienna



Elias Adriaenssens

Sascha Martens lab, University of Vienna, Max Perutz Labs - ...

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.5jyl82by7l2w/v1

Protocol Citation: Elias Adriaenssens 2024. Kinase activity assays Src and CK2. **protocols.io**

<https://dx.doi.org/10.17504/protocols.io.5jyl82by7l2w/v1>

License: This is an open access protocol distributed under the terms of the **[Creative Commons Attribution License](#)**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: July 04, 2024

Last Modified: August 29, 2024

Protocol Integer ID: 103052

Keywords: ASAPCRN

Funders Acknowledgement:

**Aligning Science Across
Parkinson's (ASAP)**

Grant ID: ASAP-000350













**Marie Skłodowska-Curie
MSCA Postdoctoral
fellowship**



Grant ID: 101062916

Abstract

This protocol details about the Kinase activity assays of Src and CK2.



- 1 To verify the activity of kinases SRC and CK2, add  45 μL of mixes containing either only kinase assay buffer ([M] 25 millimolar (mM) Tris-HCl  7.4 , [M] 150 millimolar (mM) NaCl, [M] 1 millimolar (mM) DTT, and [M] 2 millimolar (mM) MgCl_2), kinase buffer and substrate ([M] 0.5 mg/mL) or kinase buffer, substrate ([M] 0.5 mg/mL) and kinase ([M] 100 nanomolar (nM)) to individual wells of a Pierce white opaque 96-well plate (Thermo Scientific). 
- 2 Substrate peptides used were RRRDDDSDDD 10-mer (PEP-CK2I-025, Biaffin) and Poly-(Glu,Tyr 4:1) (40217, BPS) for CK2 and Src kinases, respectively. For CK2, add a specific inhibitor Silmitasertib CX-4945 (S2248, Selleck- chem), where indicated, at a concentration of [M] 1 micromolar (μM) . 
- 3 Start the reactions by the addition of  5 μL ATP in kinase assay buffer, resulting in a final concentration of [M] 100 micromolar (μM) ATP in each of the  50 μL reactions. 
- 4 After  01:00:00 at  Room temperature (RT) in darkness, add  50 μL of Kinase-Glo Max reagent (Promega) to each well, to reach a total volume of  100 μL . 

1h
- 5 Allow the luciferase reactions to stabilize for  00:15:00 before measuring luciferase activity at a Spark Multi-Mode Microplate Reader (TECAN). The luciferase activity correlates with ATP quantity, and thus, an inverse relationship between measured luminescence and kinase activity exists. 

15m