



Apr 20, 2020

Detection of SARS-Cov2 Without High Demand Reagents (Singleplex Assays)

Joseph Patterson¹, Allyson Cole-Strauss¹, John Beck¹, Caryl Sortwell¹, Jack Lipton¹

¹Michigan State University

1 Works for me dx.doi.org/10.17504/protocols.io.be8sjhwe

 Joseph Patterson
Michigan State University 

ABSTRACT

In the United States, access to testing for the novel coronavirus (SARS-Cov2) is severely limited. Arguably, the PCR based tests are the most reliable when it comes to detecting the virus. Critical reagents for these tests, however, are in short supply. Our group has worked to identify and test alternative reagents and supplies that are not in high demand by clinical labs. We have adapted a more traditional approach to isolating RNA does not use a kit. RNA isolated can be used in traditional quantitative PCR or droplet digital PCR, which has been shown to be ~500 times more sensitive.

GUIDELINES

Samples should be processed for RNA extraction (at least up until they can be frozen at -80 °C) within 48 hours of collection.

SAFETY WARNINGS

Human samples should be handled with care, and sample preparation performed in at least a BSL-2 lab.

Files



Modified Nasal Swab For Detection of Sars-Cov2
by Joseph Patterson,
Michigan State University



Isolation of SARS-Cov2 RNA from Humans Without High Demand Reagents
by Joseph Patterson,
Michigan State University



Detection of Sars-Cov2 Using Droplet Digital PCR
by Joseph Patterson,
Michigan State University



Detection of Sars-Cov2 Using qPCR
by Joseph Patterson,
Michigan State University



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits

