



Jul 14, 2022

# Expression of MicroRNA-34c in different grades of Oral squamous cell carcinoma versus normal oral mucosa (Ex-Vivo Study)

Marwa Refaat Mahmoud<sup>1</sup>, Shaymaa Omar Zayed<sup>2</sup>, Shaimaa Ali Hamouda<sup>3</sup>

<sup>1</sup>Master candidate at oral and maxillofacial pathology department , faculty of dentistry Cairo university;

<sup>2</sup>Assistant professor of oral and maxillofacial pathology,faculty of dentistry cairo university;

<sup>3</sup>lecturer of oral and maxillofacial pathology ,faculty of dentistry cairo university

1 Works for me

Share

[dx.doi.org/10.17504/protocols.io.bwjspcne](https://dx.doi.org/10.17504/protocols.io.bwjspcne)

marwa

**M** Marwa Refaat Mahmoud

## DISCLAIMER

### DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to [protocols.io](https://protocols.io) is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with [protocols.io](https://protocols.io), can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

#### ABSTRACT

The relevance of miRNAs in cancer was related to the changes in expression patterns and recurrent amplification as well as deletion of miRNA genes in cancers. miR-34c is an important tumor suppressor gene seen to be downregulated in various cancer types, such as neuroblastoma breast cancer, lung cancer and in laryngeal SCC. Previous studies investigating the expression of miR-34c in plasma and tumor tissue in patients, revealed that the high expression of miR-34c was associated with prolonged overall survival and disease-free survival compared with low expression, and it showed the possibility of regarding miR-34c as potential prognostic markers in several cancers. But to date little is known about the dysregulation of miR-34c in different grades of oral squamous cell carcinoma. The aim of our study is to evaluate the change of expression level of MicroRNA- 34c in different grades of OSCC versus normal oral mucosal tissues.

#### ATTACHMENTS

[protocol final version \(2\)  
\(4\).pdf](#)

#### DOI

[dx.doi.org/10.17504/protocols.io.bwjspcne](https://dx.doi.org/10.17504/protocols.io.bwjspcne)

#### PROTOCOL CITATION

Marwa Refaat Mahmoud, Shaymaa Omar Zayed, Shaimaa Ali Hamouda 2022.  
Expression of MicroRNA-34c in different grades of Oral squamous cell carcinoma  
versus normal oral mucosa (Ex-Vivo Study). **protocols.io**  
<https://dx.doi.org/10.17504/protocols.io.bwjspcne>



#### FUNDERS ACKNOWLEDGEMENT

#### LICENSE

\_\_\_\_\_ This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

#### CREATED

Jul 13, 2021

#### LAST MODIFIED

Jul 14, 2022

PROTOCOL INTEGER ID

51538