

JAN 19, 2024

OPEN ACCESS



DOI:

dx.doi.org/10.17504/protocols.io. 5qpvo34k9v4o/v1

Protocol Citation: Ian N Krout, Alexandria White, Tim Sampson 2024. Pole Test Assessment. protocols.io

https://dx.doi.org/10.17504/protoc ols.io.5qpvo34k9v4o/v1

License: This is an open access protocol distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working We use this protocol and it's working

Created: Jan 13, 2024

Pole Test Assessment

Ian N Krout¹, Alexandria White¹, Tim Sampson¹

¹Emory University

ASAP Collaborative Research Network

Ian Krouts Folder



Ian N Krout

ABSTRACT

Pole test is a simple behavior test used to assess fine motor coordination. Mice are placed at the top of a 50-60 cm vertical pole with a diameter of 1 cm. The pole is attached to a base that is placed in the home cage so that mice might prefer to descend to the floor of cage. The test is conducted over 3 days: training day 1, training day 2, and test day. Each mouse has 3 trials during the test day as well as on training days. Daily sessions should be consecutive (i.e., do not let training and test days be separated by a weekend). Record each trial on test day with a camera to obtain accurate measures of both time-to-turn (TTT) and total-time-to-descend (TTD).

Oct 19 2024



Last Modified: Jan 19, 2024

PROTOCOL integer ID: 93488

Keywords: ASAPCRN, Behavior, Mouse, PD, Neurobehavior

Funders Acknowledgement:

Aligning Science Across Parkinson's

Grant ID: ASAP020537

MATERIALS

- 1. A metal pole with a diameter of 1cm and height of 50-60cm, this should come on a base. (See attached photo).
- 2. Foam/PVC liner, we use a cabinet liner similar to what can be found here. Adhesive Grip for Pole Test.
- 3. Standard Mouse cages.
- 4. Stopwatch/Timer.
- 5. GoPro/Means of recording each trial.




Figure 1. Example of pole test apparatus used by our lab.

Pole set-up

protocols.io


- 1 Find a metal pole 1cm in diameter with a height of ~50-60cm. This should have a base that is able to fit inside of a standard mouse cage.
- Wrap the pole with an adhesive foam or PVC material (see material for link). This should cover the entire pole and be replaced as it wears down.
- 3 Tape the top and bottom of the adhesive to secure to the pole.
- 4 Wipe down the base and pole with 70% EtOH before beginning any trials and between each mouse.

Acclimation

On each training and assessment day bring the mice up from the vivarium to the behavior room at least 1h prior to assessment to allow for acclimation to the environment prior to assessment.

Training Day 1

- **6** Determine the order of assessment that you will use for the length of the training and assessment, record this in lab notebook.
- Remove all mice from the first cage assessed and place into a clean empty mouse cage. Pole test training and assessment will be performed in the home cage.



Training Day 2

Repeat all steps from training day 1 no more than 24h later. At this point the mice should be able to perform all trials without issue.

Assessment

- No more than 24h after training day 2 the mice should be assessed for their ability to descend the pole.
- **15** Repeat steps 6-9 for each mouse.
- 16 Start recording, so that each mouses trial is captured.
- Place the mouse face up on top of the pole, and this time do not intervene with your hand. By now the mouse should not attempt to go over the pole. If it does, begin the trial again.
- Begin the stopwatch/timer as soon as you place the mouse on the pole. Stop timing once it fully descends and both front paws are touching the ground.
- Record the time-to-descend (TTD) in your notebook.

20	Allow the mouse to free roam post descent for 30 seconds.
21	Repeat this 2 more times (total of 3 trials) for each mouse.
22	Clean the pole and base between each mouse using 70% EtOH.
23	After all mice in a cage are assessed, replace mice in the cage and place pole (cleaned) into a new home cage.
	Analysis
24	Two measures are analyzed for this pole test assessment.
25	Time to decond (TTD) is calculated by taking the average time to decond the male frame the 2 trials of each
25	Time-to-descend (TTD) is calculated by taking the average time to descend the pole from the 3 trials of each mouse. This measure can be compared between genotypes, treatment groups, cages, etc.
25 26	

- 27.1 Mouse won't or cannot turn around and/or descend = Physically Unable to Perform (PUP)
- 27.2 Mouse falls off or slides down pole (shouldn't happen if pole is wrapped in grippy material, such as electric tape or ideally non-slip shelf liner) = PUP
- 27.3 Mouse descends sideways and never fully aligns its body with pole = count as TTT, no descent time recorded.
- Mouse nearly descends, but goes back up the pole again = PUP/exclude trial