




NOV 11, 2022

 SHARE

WORKS FOR ME 1


 Comparison of measured LDL cholesterol with calculated LDL-cholesterol using the Friedewald and Martin-Hopkins formulae in diabetic adults at Charlotte Maxeke Johannesburg Academic Hospital/NHLS Laboratory

DOI

dx.doi.org/10.17504/protocols.io.yxmvm2ky6g3p/v1

Mogomotsi Dintshi¹, Ngalulawa Kone¹, Siyabonga Khoza¹

¹Departement of Chemical Pathology, National Health Laboratory Services and University of Witwatersrand, Johannesburg, South Africa.



Mogomotsi Dintshi

COMMENTS 0

ABSTRACT

Background

National Cholesterol Education Programme Adult Treatment Panel III (NCEP ATP III) and the European Society of Cardiology recommends using low-density lipoprotein cholesterol (LDL-C) as a treatment target for cholesterol lowering therapy. The Friedewald formula underestimate LDL-C in non-fasted and hypertriglyceridemia patients. This study aimed to compare measured LDL-C to calculated LDL-C in diabetic patients using the Friedewald and Martin-Hopkins formulae.

Methods

The data of 1 247 adult diabetes patients were retrospectively evaluated, and included triglycerides (TG), LDL-C, total cholesterol, and high-density lipoprotein cholesterol that were measured on the Roche Cobas® c702. Passing-Bablok regression analysis was used to determine the degree of agreement between measured LDL-C and calculated LDL-C using both formulae. The Bland-Altman plots were used to assess the bias at medical decision limits based on the 2021 European Society of Cardiology (ESC) guidelines on cardiovascular disease prevention in clinical practice.

Results

Both formulae showed a good linear relationship against measured LDL-C. However, the Martin-Hopkins formula outperformed the Friedewald formula at LDL-C treatment target $<1.4\text{mmol/L}$. The Friedewald formula and the Martin-Hopkins formula had 14.9% and 10.9% mean positive bias, respectively. At $\text{TG-C} \geq 1.7\text{ mmol/L}$, the Martin-Hopkins formula had a lower mean positive bias of 4.2 % (95 % CI 3.0-5.5) compared to the Friedewald formula, which had a mean positive bias of 21.8 % (95 % CI 19.9-23), which was higher than the NCEP ATP III recommended total allowable limit of 12%.

Conclusion

The Martin-Hopkins formula performed better than the Friedewald formula at LDL-C of 1.4 mmol/L and showed the least positive bias in patients with hypertriglyceridemia.

DOI

dx.doi.org/10.17504/protocols.io.yxmvm2ky6g3p/v1

PROTOCOL CITATION

Mogomotsi Dintshi, Ngululawa Kone, Siyabonga Khoza 2022. Comparison of measured LDL cholesterol with calculated LDL-cholesterol using the Friedewald and Martin-Hopkins formulae in diabetic adults at Charlotte Maxeke Johannesburg Academic Hospital/NHLS Laboratory. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.yxmvm2ky6g3p/v1>



KEYWORDS

Calculated low density lipoprotein; Diabetes mellitus; Direct low density lipoprotein; Friedewald formula; Martin-Hopkins formula

LICENSE

_____ This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Oct 10, 2022

LAST MODIFIED

Nov 11, 2022

PROTOCOL INTEGER ID

71084

MATERIALS TEXT

 Research report data (1).xlsx