



NOV 28, 2022

WORKS FOR ME

1



## MycoFluor Mycoplasma Detection

DOI

[dx.doi.org/10.17504/protocols.io.bp2l692nklqe/v1](https://dx.doi.org/10.17504/protocols.io.bp2l692nklqe/v1)Ali Albalakhi<sup>1,2</sup>, Ning Xia<sup>1,2</sup><sup>1</sup>Massachusetts General Hospital;<sup>2</sup>Aligning Science Across Parkinson's (ASAP) Collaborative Research Network

COMMENTS 0



Daniel Choo

## ABSTRACT

MycoFluor™ Mycoplasma Detection produces a fast and simple fluorescence microscopic assay that identifies mycoplasma infection in cell cultures. In order to detect mycoplasma, the fluorescent MycoFluor™ reagent is added to the culture medium, with or without cells, and the sample becomes stained and examined under a fluorescence microscope.

DOI

[dx.doi.org/10.17504/protocols.io.bp2l692nklqe/v1](https://dx.doi.org/10.17504/protocols.io.bp2l692nklqe/v1)

## PROTOCOL CITATION

Ali Albalakhi, Ning Xia 2022. MycoFluor Mycoplasma Detection . **protocols.io**  
<https://dx.doi.org/10.17504/protocols.io.bp2l692nklqe/v1>



## LICENSE

———— This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

## CREATED

Nov 23, 2022

## LAST MODIFIED

Nov 28, 2022

## PROTOCOL INTEGER ID

73200

## MATERIALS TEXT

## Reagents Needed:

1. MycoFluor Mycoplasma Detection Kit (M-7006)
2. Microcentrifuge tubes
3. Microscope slides
4. Clear coverslips

## Protocol Testing of Culture Media

- 1 Take about 4mL of cell medium directly from the culture dish in which the cells have been growing centrifuge the sample at 1300 x g for 10min to pellet any cells and debris
- 2 Carefully transfer 1mL of supernatant into labeled microcentrifuge tubes.
- 3 Centrifuge the microcentrifuge tubes at 12,500 x g for 15 minutes
- 4 Carefully remove and discard 0.5mL of supernatant, leaving behind 0.5mL of medium in the tube. Resuspend any pellet that may have formed using this 0.5mL of medium.
- 5 Add 26µL of 20X concentrated MycoFluor reagent to 0.5mL of the medium.
- 6 Pipet 10µL of the stained medium onto a clean microscope slide and cover with a clean coverslip.
- 7 Seal the slide using quick dry clear nail polish topcoat by covering all sides of the coverslip.
- 8 Image the slide using fluorescent microscope.

## Control Slides with Mycoplasma MORFS

- 9 Generating positive controls for the testing of culture media

- 9.1 a. Pipet 5 $\mu$ L of the mycoplasma MORFS stock suspension on a clean, labeled microscope slide
- 9.2 b. Add 5 $\mu$ L of stained medium to the slide
- 9.3 c. Cover with clear coverslip and seal using quick dry topcoat nail polish.
- 10 Image the positive controls and compare to the samples that have not been spiked with MORFS

## Microscopy

- 11 Prepare the microscope with a near ultraviolet fluorescence filter (Excitation at 365nm and either bandpass 450nm  $\pm$  30nm or longpass >400nm emission filter)
- 12 For optimum result, a 100X oil immersion objective is suggested but a 60X oil immersion objective will also work
- 13 First look at the control slide spiked with MORFS to get an idea what it should look like. Then examine the test slide for extranuclear blue fluorescence.