

Transforming the marine bacterium *Ruegeria pomeroyi* using tri-parental mating

APR 10, 2023

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.yxmvmk6o6g3p/v1

Protocol Citation: Cherry Gao, Christopher R. Reisch, Mary Ann Moran, Roman Stocker 2023. Transforming the marine bacterium *Ruegeria pomeroyi* using tri-parental mating. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.yxmvmk6o6g3p/v1>

MANUSCRIPT CITATION:
 Gao, C., Fernandez, V.I., Lee, K.S. et al. Single-cell bacterial transcription measurements reveal the importance of dimethylsulfoniopropionate (DMSP) hotspots in ocean sulfur cycling. *Nat Commun* 11, 1942 (2020).
<https://doi.org/10.1038/s41467-020-15693-z>
 Gao, C. Ecological insights through single-cell measurements of marine bacteria. Ph. D. Thesis, Massachusetts Institute of Technology, Department of Biological Engineering, September 2020.
<https://hdl.handle.net/1721.1/129213>

Transforming the marine bacterium *Ruegeria pomeroyi* using tri-parental mating

Cherry Gao^{1,2,3}, Christopher R. Reisch⁴, Mary Ann Moran⁵, Roman Stocker³

¹Department of Biological Engineering, Massachusetts Institute of Technology, Cambridge, MA 02139, USA;


²Department of Civil and Environmental Engineering, Ralph M. Parsons Laboratory, Massachusetts Institute of Technology, Cambridge, MA 02139, USA;

³Department of Civil, Environmental and Geomatic Engineering, Institute for Environmental Engineering, ETH Zurich, 8093 Zurich, Switzerland;

⁴Department of Microbiology and Cell Science, University of Florida, Gainesville, FL 32611, USA;

⁵Department of Marine Sciences, University of Georgia, Athens, GA, USA

Environmental Microfluidics Laboratory



Cherry Gao

DISCLAIMER

Please reference the following publication:

Gao et al., Single-cell bacterial transcription measurements reveal the importance of dimethylsulfoniopropionate (DMSP) hotspots in ocean sulfur cycling. *Nature Communications*. 11, 1942 (2020).
doi.org/10.1038/s41467-020-15693-z

ABSTRACT

This protocol describes a tri-parental mating method for transforming the model marine bacterium *Ruegeria pomeroyi* DSS-3 with a desired recombinant plasmid. This transformation method is reliable, easy-to-use, and does not require specialized equipment. Using this method, we have successfully transformed *R. pomeroyi* with large plasmids (up to 14.833 kb) which enabled the construction of the first fluorescent reporter strains of *R. pomeroyi* (described in doi.org/10.1038/s41467-020-15693-z).

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Sep 06, 2021

Last Modified: Apr 10, 2023

PROTOCOL integer ID:
53012

Keywords: Ruegeria pomeroyi, genetic engineering, cloning, transformation, triparental mating, fluorescence reporter

ATTACHMENTS

[Transforming the marine bacterium Ruegeria pomeroyi using tri-parental mating.pdf](#)

MATERIALS

Materials:

- E. coli donor strain, containing the plasmid to be transformed into R. pomeroyi
- E. coli helper strain, containing the helper plasmid pRK600 plasmid (ampicillin resistant)
- R. pomeroyi, wildtype
- 1/2 YTSS liquid medium (in 500 ml⁻¹ : 2 g yeast extract (BD Biosciences), 1.25 g tryptone (BD Biosciences), 10 g sea salts (Sigma-Aldrich))
- 1/2 YTSS agar plate (in addition to 500 ml liquid medium, 7.5 g agar (Bacto Agar, BD Biosciences)), pre-warmed to 30 °C
- Luria Broth (LB) liquid medium
- Antibiotics kanamycin and chloramphenicol
- Potassium tellurite (Fluorochem)
- Sterile L-shape cell spreader
- Parafilm
- Eppendorf tubes (1.5 ml; sterile)
- Incubators (with shakers) at 30 °C and 37 °C
- Table-top centrifuge

SAFETY WARNINGS



Potassium tellurite is used to select for R. pomeroyi, and against E. coli. Potassium tellurite is toxic if swallowed, and causes skin and serious eye irritation. Furthermore, it is possible that R. pomeroyi mediates the formation of volatile tellurides which may be highly toxic. Thus, care should be used to avoid contact with the substance, and colony picking may best be performed in a fume hood to protect against the potential presence of volatile tellurides.



BEFORE START INSTRUCTIONS

Before using this protocol, the user should have designed and built the plasmid desired for transforming into *R. pomeroyi*. The authors have built and successfully transformed reporter plasmids on vector backbones pBBR1MCS-KanR (GenBank U23751) and pRK415 (GenBank EF437940). Furthermore, if the desired plasmid is small (empirically, the authors suggest less than 11 kb), electroporation may be a faster method for transforming *R. pomeroyi*. Recommendations for plasmid design and electroporation method for *R. pomeroyi* are available in detail in Cherry Gao's PhD Thesis (<https://hdl.handle.net/1721.1/129213>).


Execution



5m



1

Prepare  Overnight  5 mL liquid cultures of the three bacterial strains for tri-parental mating:



(1) wildtype *R. pomeroyi* (1/2 YTSS medium,  30 °C with shaking);

(2) helper *E. coli* containing the pRK600 plasmid (LB medium amended with  15 µg/mL chloramphenicol ,  37 °C with shaking)




(3) donor *E. coli* containing the constructed plasmid (LB medium amended with  50 µg/mL kanamycin ,  37 °C with shaking).

2

After overnight growth, wash the two *E. coli* cultures twice in 1/2 YTSS medium to eliminate antibiotics.




3

To wash, aliquot  1 mL overnight cultures into Eppendorf tubes, pellet the bacterial cells through gentle centrifugation  3000 rpm, 00:05:00 (on a tabletop centrifuge), discard the supernatant, and resuspend in  1 mL 1/2 YTSS medium .





5m

4


Pellet the *E. coli* cells again through centrifugation, discard the supernatant, then resuspend in  1 mL YTSS medium .




5

In a clean Eppendorf tube, combine the following:  2 mL overnight culture *R. pomeroyi* ,  200 µL washed overnight culture helper *E. coli* and

 200 µL washed overnight culture of donor *E. coli* .



- 6 Gently centrifuge this mixture to pellet the bacterial cells. Discard the supernatant and resuspend in  100 µL 1/2 YTSS medium .



- 7 Deposit  50 µL concentrated bacterial mixture onto the center of a pre-warmed 1/2 YTSS plate. Place the lid onto the plate.


- 8 Incubate the plate (lid-side up)  Overnight at  30 °C to allow mating to occur.



- 9 For selection of plasmid-containing *R. pomeroyi*, pick up most of the biomass (now dry) that is at the center of the mating plate with a sterile L-shaped cell spreader, and evenly spread this biomass onto a 1/2 YTSS plate amended with  50 µg/mL kanamycin and  50 µg/mL potassium tellurite .


Note

Kanamycin and potassium tellurite stock solutions (1000× concentration) should be prepared by dissolving desired amounts in milliQ water and filter sterilizing (0.2 µm; not autoclaved). Stock solutions can be added to melted 1/2 YTSS agar medium before pouring into plates, or spread directly onto solidified 1/2 YTSS agar plates. To directly spread onto solidified agar plates, place ~20 µl of the 1000x stock solution in the middle of the plate and spread evenly using a sterile L-shaped cell spreader. Let the compounds diffuse through the agar medium for at least 1 hour at room temperature before applying bacteria.

- 10 Incubate the selection plate (lid-side down) at  30 °C until colonies of transformed *R. pomeroyi* form. Typically, small colonies become visible within 48–72 hours. Due to the presence of potassium tellurite, *R. pomeroyi* colonies appear dark brown.



- 11 To avoid evaporation during this long incubation time, secure the lid to the plate with Parafilm.

- 
- 12 Confirm successfully transformed *R. pomeroyi* through colony PCR and sequencing. If possible, 16S sequencing to confirm the identity of *R. pomeroyi* is recommended.