



2 ▼

Apr 01, 2022

Design and preparation of synthetic reference peptides for APP/A β TOMAHAQ proteomics V.2

Hankum Park^{1,2,3}, Frances V Hundley^{1,2}, Harper JW^{1,2}¹Department of Cell Biology, Harvard Medical School Boston, MA 02115, USA;²Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD 20815, USA;³Current affiliation: Soeul National University, School of Dentistry

1

dx.doi.org/10.17504/protocols.io.bp2l6bqk1gqe/v2

Frances Hundley

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to protocols.io is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with protocols.io, can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

TOMAHQA-based targeted proteomics relies on heavy-labeled reference peptides for multi-plexed quantification of peptides of interest within a set of samples. The APP amyloid precursor protein is thought to be proteolytically processed within the endolysosomal system by β -secretase and γ -secretase to yield various forms of A β . To quantitatively track APP products, we describe a protocol for design and generation of synthetic reference peptides for TOMAHQA analysis. We also include reference peptides for the extracellular and cytosolic domains.

[synthetic peptides for
protocols.io.xlsx](#)

DOI

dx.doi.org/10.17504/protocols.io.bp2l6bqk1gqe/v2

Hankum Park, Frances V Hundley, Harper JW 2022. Design and preparation of synthetic reference peptides for APP/A β TOMAHAQ proteomics. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bp2l6bqk1gqe/v2>
Frances Hundley



ASAPCRN

_____ protocol ,

Mar 25, 2022

Apr 01, 2022

59928

:

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to [protocols.io](https://dx.doi.org/10.17504/protocols.io.bp2l6bqk1gqe/v2) is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with [protocols.io](https://dx.doi.org/10.17504/protocols.io.bp2l6bqk1gqe/v2), can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

- 1 Design APP peptides corresponding to extracellular and cytosolic regions based on data available in Peptide Atlas which are predicted to have favorable LC-MS properties. See attached Table.
- 2 Design half-tryptic peptides based on the cleavage sites for BACE1 and γ -secretase. See attached table.
- 3 Order commercial synthesis of peptides from Biomatik and Thermo Fischer Scientific, or similar.

- 4 Reconstitute the peptides with 3% acetonitrile/0.1% formic acid, and quantify the concentration using Pierce Quantitative Fluorometric Peptide Assay.
- 5 Mix 10 μL of 200 μM peptide in 200 mM EPPS buffer (pH 8.5) with 1 μL of 5 $\mu\text{g}/\mu\text{L}$ super-heavy TMT reagent (TMTsh). Incubate at 25 °C for 1 hr then check the labeling efficiency by LC-MS.
- 6 Add extra TMTsh to under-labeled peptides until > 95% labeling on both N-terminus and lysine residues is reached.
- 7 Desalt the reaction mixture using C18 StageTip.
- 8 Resuspend the peptides in 5% acetonitrile/8% formic acid, and pool the peptides.